

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2001 (10.05.2001)


PCT

(10) International Publication Number
WO 01/32711 A2

- (51) International Patent Classification⁷: C07K 14/75
- (21) International Application Number: PCT/US00/29187
- (22) International Filing Date: 23 October 2000 (23.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/160,608 21 October 1999 (21.10.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/160,608 (CIP)
Filed on 21 October 1999 (21.10.1999)
- (71) Applicant (for all designated States except US):
BOARD OF TRUSTEES OF THE UNIVERSITY
OF ARKANSAS [US/US]; 2404 North University Av-
enue, Little Rock, AR 72207-3608 (US).
- (72) Inventor; and
(75) Inventor/Applicant (for US only): HERMONAT, Paul,
L. [US/US]; 1812 Reservoir Road, Apartment 143, Little
Rock, AR 72227 (US).
- (74) Agent: GRANADOS, Patricia, D.; Foley & Lard-
ner, 3000 K. Street, N.W., Suite 500, Washington, DC
20007-5109 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,

[Continued on next page]

(54) Title: RepO-ASSOCIATED VIRUS AAV REP78 MAJOR REGULATORY PROTEIN, MUTANTS THEREOF AND USES THEREOF



(57) Abstract: AAV Rep78 mutants comprising a modified AAV Rep78 protein that possess different biochemical and biological functions as compared to the wild-type AAV Rep78 protein are disclosed. Particularly, the AAV Rep78 mutants that bind to at least one of a papillomavirus DNA or an AAV DNA or an oncogene or HIV DNA differently as compared to the wild-type AAV Rep78 mutants are

[Continued on next page]

WO 01/32711 A2



NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

Published:

— Without international search report and to be republished
upon receipt of that report.

(84) **Designated States (regional):** ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

disclosed. The present invention further discloses DNA sequences encoding at least one AAV Rep 78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein and pharmaceutical compositions comprising the DNA sequences. The present invention additionally is directed to a method of inhibiting papillomavirus associated diseases comprising administering pharmaceutical compositions containing AAV Rep78 mutants or the DNA sequences encoding the mutants. Also disclosed are DNA sequences comprising the full length AAV genome modified to where the AAV Rep78 protein is replaced with an AAV Rep mutant that binds weakly or not at all to a papillomavirus DNA or an AAV DNA or both differently as compared to the wild-type AAV Rep78 protein and its use in producing recombinant AAV at increased levels production over wild-type levels of production. Further disclosed is a method of inhibiting papillomavirus-associated diseases, cancer, and HIV-associated diseases comprising administering a pharmaceutical composition comprising a wild-type AAV Rep78 or a mutant thereof to a patient afflicted with a papillomavirus-associated disease, cancer, or HIV-associated diseases. Particularly, useful in treating papillomavirus-associated diseases are the AAV Rep78 or mutant thereof that binds to nucleotides 14-56 of p97 of HPV-16 and inhibits expression of HPV oncoproteins. An AAV Rep78 regulation element of a DNA sequence comprising about nucleotides 14-56 of the nucleotide sequence of Figure 2 and an AAV Rep78 regulatable promoter comprising the regulation element and a promoter sequence except for the HPV-16 p97 promoter.

RepO-ASSOCIATED VIRUS AAV REP78 MAJOR REGULATORY PROTEIN, MUTANTS THEREOF AND USES THEREOF

5

BACKGROUND OF THE INVENTION

The present invention is directed to adeno-associated virus (AAV) Rep78 mutants and DNA sequences encoding these mutants that possess the capability (a) to inhibit papillomavirus (PV) and PV-associated diseases or cancer or human immunodeficiency virus (HIV) and HIV-associated diseases or (b) to enhance AAV functions and DNA sequences encoding these AAV Rep78 mutants. The present invention also is directed to AAV Rep78 mutants and DNA sequences encoding these mutants that possess the capability to inhibit PV transcriptional activity, or alternatively, possess the capability to enhance AAV functions and the DNA constructs containing these DNA sequences. The present invention is directed to a method of inhibiting PV and PV-associated diseases or cancer or HIV and HIV-associated diseases comprising administering a composition comprising the AAV Rep78 mutants or the wild-type AAV Rep78 major regulatory protein (AAV Rep78 protein) or the DNA sequences encoding these mutants or proteins to a patient having a PV-associated disease or cancer or HIV-associated diseases. The present invention further is directed to DNA constructs encoding an AAV Rep78 protein or mutant thereof, operably linked to a specific inducible promoter that is induced during PV infection or cancer or HIV infection and the DNA construct's use in inhibiting PV-associated diseases or cancer or HIV-associated diseases in a patient. The present invention also provides a method for producing increased amounts of recombinant AAV (rAAV). A specific regulation element specific for AAV Rep78 binding is disclosed.

25 The present invention is further directed to a method of improving the level of rAAV gene therapy and pseudo-wild-type AAV production by all forms of complementor/generator systems by substituting improved, non-inhibiting AAV Rep78 mutants in place of the wild-type AAV Rep78 protein. There are many techniques for generating rAAV particles for gene therapy. These techniques are improved by using a mutant AAV Rep78, such as AAV Rep77^{LG} and AAV Rep79^{FA} genes or proteins, and other AAV Rep78 mutants that possess comparable properties, disclosed in this application that bind weakly or not at all to DNA sequences from PV, AAV, oncogenes or HIV,

30

particularly, HIV-1 or a combination thereof, and preferably promoter sequences from these sources. The designations of these mutants indicate that the number following "Rep" is the amino acid that is changed. The specific amino acid change is shown in the superscript where the first amino acid is the amino acid in the wild-type AAV Rep78 and the second amino acid is the amino acid that the wild-type amino acid is changed to in the AAV Rep mutant. For example, for AAV Rep77^{LG}, the amino acid 77 is leucine (L) in the wild-type AAV Rep 78 and it is changed to glutamine (G) in the AAV Rep mutant. These designations apply to AAV Rep78 mutants disclosed in the present invention. These proteins have all of the activities required for the AAV life cycle but do not have the same ability to inhibit transcription that the wild-type AAV Rep78 protein does. Thus, AAV production is increased by 100-200% when using a vector containing one of these mutants, and therefore, these mutants are useful for producing higher titers of rAAV stock.

Other AAV Rep78 mutants that are useful within the context of the present invention, are mutants that bind better than wild-type AAV Rep78 to DNA sequences from PV, AAV, oncogenes or HIV, particularly, HIV-1, or a combination thereof, and preferably promoter sequences from these sources. AAV Rep 192^{HG}, is an example of one of these mutants, that binds to AAV p5 promoter and AAV TR and inhibits HPV-16 oncogenic transformation better than wild-type AAV Rep78. AAV Rep 192^{HG} and other AAV Rep78 mutants with comparable binding affinities are useful inhibitors of HPV and AAV promoters, and therefore, useful in inhibiting PV and PV-associated diseases, as well as HIV and HIV-associated diseases. These mutants also are useful in anti-HIV protocols and in anti-cancer protocols.

Infection and DNA integration by certain HPV types are central events in the generation of cervical cancer (Durst, *et al.*, *Proc. Natl. Acad. Sci. USA* 80:3812-3815 (1983), Cullen, *et al.* *J. Virol.* 65:606-612 (1991)). The most common HPV type associated with cervical cancer is HPV-16, and roughly two thirds of cervical cancers contain the DNA of this virus. The HPV-16 sequence is provided in Seedorf *et al.*, *Virology* 145(1):181, (1985). AAV is another virus commonly found in the anogenital region (Blacklow, *et al.*, *Proc. Natl. Acad. Sci. USA* 58:1410-1415 (1967), Bantel-Schaal, *et al.*, *Virology* 134:52-63 (1984), Tobiasch, *et al.*, *J. Med. Virol.* 44:215-222 (1995), Han, *et al.*, *Virus Genes* 12:47-52 (1996)). AAV is described as a helper dependent human parvovirus, which requires the cell to be co-infected with an adenovirus, herpes virus or

pox virus, for AAV DNA replication and virus particle formation to take place. Without these helper viruses, AAV is unable to replicate, except under special circumstances. Adenovirus is believed to be the most common helper virus *in vivo*. When no helper virus is present, AAV will set up a latent infection by chromosomal integration, specifically in a region on chromosome 19.

Infection by AAV, in sharp contrast to the HPV, is negatively associated with cervical cancer as demonstrated by the prevalence or titer of anti-AAV antibodies (Mayor, *et al.*, *Am. J. Obstet. Gynecol.* 126:100-105 (1976), Georg-Fries, *et al.*, *Virology* 134:64-71 (1984)). Bidirectional interaction has been observed between AAV and PVs. One such interaction is that PVs might serve as helpers for AAV (Walz, *et al.*, *J. Gen. Virol.* 78:1441-1452 (1997)), allowing for AAV DNA replication and virion production.

In addition to its negative association with PV infections, AAV is a useful viral vector for human gene therapy. The nucleotide sequence and organization of the AAV 2 genome is known and published. (Srivastava *et al.*, *J. Virol.* 45 (1) 555, (1983)). AAV latently infects cells as a natural part of its life cycle. AAV will latently infect the cell by chromosomal integration, usually within a 100 base pair region on human chromosome 19. Furthermore, AAV has not been found to be the etiologic agent of any disease, and is thus accurately described as being non-pathogenic. These properties suggest that AAV is an ideal gene transmitter for human gene therapy. AAV is highly effective in transducing a variety of cell types in tissue culture including cells of epithelial, fibroblastic and hematopoietic origin.

AAV Rep78 is an AAV DNA binding transcription factor and was first identified in Mendleson *et al.*, *J. Virol.* 60(3) 823 (1986), as the largest of four products encoded by the AAV *rep* open reading frame. AAV Rep78 is required for AAV DNA replication (Hermonat *et al.*, *J. Virol.* 51: 329, (1984); Tratschin *et al.*, *J. Virol.* 51: 611, (1984)), for AAV gene regulation (Labow *et al.*, *J. Virol.* 60: 251, (1986); Tratschin *et al.*, *Mol. Cell. Biol.* 5: 3251, (1986)), and AAV Rep78 displays a variety of biochemical activities which are necessary for these biological phenotypes (Im *et al.*, *Cell.* 61: 447, (1990); Ni *et al.*, *J. Virol.* 68: 1128, (1994)).

AAV Rep78 is the full length product of the *rep* (for replication) open reading frame. The AAV Rep78 protein is critical for the AAV life cycle. In 1984, the initial genetic analysis of AAV using a series of linker insertion-frame shift mutants was carried

out (Hermonat *et al.*, *J. Virol.* 51: 329, (1984)). In this study, the *rep*, *lip*, and *cap* mutant phenotypes were discovered and the corresponding genes within AAV were mapped. The data from this study shows that mutations at map units 11, 32, and 42 were defective for DNA replication. The positioning of these mutations map the effect to the AAV Rep78 protein, as only AAV Rep78 is affected by all three mutations. There are three other Rep proteins, Rep68, Rep52, and Rep40, which are produced from the same open reading frame (ORF), and are largely subsets of the AAV Rep78 amino acid sequence. It is unclear if these proteins have any unique biochemical properties. They may serve to augment some of AAV Rep78's functions. In any case, AAV Rep78 is all that is needed for the AAV life cycle as indicated by earlier genetic studies and by recent data in which only AAV Rep78, of the four Rep proteins, was needed for generating rAAV vectors (Holscher *et al.*, *J. Virol.* 69: 6880 (1995)).

AAV Rep78 interacts with DNA. One of AAV Rep78's functions is to bind to the AAV terminal repeats (TR DNA) which functions as the origin of replication (Ryan *et al.*, *J. Virol.* 70:1542 (1996); McCarty *et al.*, *J. Virol.* 68:4988 (1994); McCarty(1996) *et al.*, *J. Virol.* 68:4998 (1994); Snyder *et al.*, *J. Virol.* 67:6096 (1993); Ashktorab *et al.*, *J. Virol.* 63:3034 (1989)).

In earlier studies, it has been demonstrated that AAV inhibits bovine papillomavirus type 1 (BPV-1) and HPV-16-induced oncogenic transformation (Hermonat, P.L., *Virology* 172:253-261 (1989); Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994); Hermonat, *et al.*, *Gynecologic Oncology* 66:487-494 (1997)). Others have also observed AAV inhibition of BPV-1 and HPV-18 (Schmitt, *et al.*, *Virology* 172:73-81 (1989); Horer, *et al.*, *J. Virol.* 69:5485-5496 (1995); Su, *et al.*, *British J. Can.* 73:1533-1537 (1996)) and HIV-1. The effect has been mapped to the AAV encoded Rep78 protein, and this protein has been shown to inhibit expression of the PV promoter just upstream of the E6 gene (p89 of BPV-1, p97 of HPV-16, and p105 of HPV-18) (Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994); Schmitt, *et al.*, *Virology* 172:73-81 (1989); Horer, *et al.*, *J. Virol.* 69:5485-5496 (1995); Hermonat, *et al.*, *Virology* 247(2):240-250 (1998)). Similar to PV, HIV has been shown to be inhibited by AAV Rep78 (Antoni BA. *et al.*, *J. Virology*. 65(1):396-404 (1991); Rittner K. *et al. Biochemical Society Transactions.* 19(4):438S (1991); Oelze I. *et al.*, *J. Virology.* 68(2):1229-33 (1994)), and DNA binding is required for this inhibition (Batchu and Hermonat, *FEBS Letters*, 367(3): 267-71 (1995); Kokorina

NA. *et al.*, *J. Human Virology* 1(7):441-50 (1998)). Rep78 binds to two DNA sequences of the HIV long terminal repeat promoter, the TAR sequences (nt +23 to +42) (Batchu and Hermonat, *FEBS Letters*, 367(3):267-71 (1995)) and to a region just upstream of the TATA box (nt -54 to -34).

- 5 AAV Rep78 regulates a variety of heterologous genes. C-H-ras (Katz, *et al.*, *Cancer Research* 46:3023-3026 (1986); Hermonat, P.L., *Cancer Research* 51:3373-3377 (1991); Khleif, *et al.*, *Virology* 181:738-741 (1991)), c-fos (Klein-Bauernschmitt, *et al.*, *J. Virol.* 66:419-4200 (1992); Hermonat, P.L., *Cancer Letters* 81:129-136 (1994)), c-myc (Klein-Bauernschmitt, *et al.*, *J. Virol.* 66:419-4200 (1992); Hermonat, P.L., *Cancer Letters* 81:129-136 (1994)), and the HIV long terminal repeat (HIV-LTR) (Rittner, *et al.*, *J. Gen. Virol.* 73:2977-2981 (1992); Antoni, *et al.*, *J. Virol.* 64:396-404 (1991)) are down-regulated by AAV Rep78, while the c-sis promoter is up-regulated (Wonderling, *et al.*, *J. Virol.* 70:4783-4786 (1996)). Still other genes are not affected, such as the murine osteosarcoma virus long terminal repeat (MSV-LTR) (Hermonat, P.L., *Cancer Research* 15 51:3373-3377 (1991)) and the human β -actin promoter (Horer, *et al.*, *J. Virol.* 69:5485-5496 (1995)). The largest of 4 products encoded by the AAV *rep* open reading frame (Mendleson, *et al.*, *J. Virol.* 60:823-832 (1986)), AAV Rep78, is required for AAV DNA replication (Hermonat, *et al.*, *J. Virol.* 51:329-333 (1984); Tratschin, *et al.*, *J. Virol.* 51:611-619 (1994)) and for AAV gene regulation (Labow, *et al.*, *J. Virol.* 60:251-258 20 (1986); Tratschin, *et al.*, *Mol. Cell. Biol.* 5:3251-3260 (1986)). AAV Rep78 carries out a range of biochemical activities which are necessary for its biological phenotypes (Im, *et al.*, *Cell* 61:447-57 (1990); Ni, *et al.*, *J. Virol.* 68:1128-1138 (1994)), including binding to promoter DNA (McCarty, *et al.*, *J. Virol.* 74:4988-4997 (1994); Batchu, *et al.*, *Cancer Letters* 86:23-31 (1994); Wonderling, *et al.*, *J. Virol.* 71:2528-2534 (1996)), and to a 25 variety of cellular proteins (Hermonat, *et al.*, *Biochem. and Molec. Biol. Internat.* 403:409-420 (1997)), including the transcription factors Sp1 (Hermonat, *et al.* *Cancer Research* 56:5299-5304 (1996); Pereira, *et al.*, *J. Virol.* 71:1747-1756 (1997)), TBP (Hermonat, *et al.*, *Virology* 245:120-127 (1998)), and itself (Weitzman, *et al.*, *J. Virol.* 70:2440-2448 (1996), Hermonat, *et al.*, *FEBS Letters* 401:180-184 (1997), Smith, *et al.*, *J. Virol.* 30 71:4461-4471 (1997)).

Hermonat *et al.*, *Cancer Research* 51: 3373 (1994) involved the use of the complete or partial genomes and did not study the p97 promoter in isolation as does the present

invention. This is underscored by the identification of a new promoter in the E7 gene (Higgins *et al.*, *J. Gen. Virol.* 73 (Pt. 8):2047, (1992)), which was included in all of the plasmids used in the Hermonat *et al.* study.

Although Hermonat *et al.*, *Gynecologic Oncology* 66: 487 (1997) suggests that
5 AAV Rep78 specifically inhibits the HPV-16 p97 promoter, this publication itself acknowledges its own weakness by stating that the study does not mutationally map the specific AAV rep region product which is required for the inhibition of oncogenic transformation. Further, this publication suggests that AAV Rep78 affects p97
10 transcription based upon results obtained with a chloramphenicol acetyltransferase (CAT) assay. This assay measures both transcription and translation as the activity measured is of the resulting protein. In fact, the ability of AAV Rep78 to affect translation, and not transcription, was emphasized in Trempe *et al.*, *J. Virol.* 62(1): 68 (1988). Thus, the main effect on p97 was not definitively proven by Hermonat *et al.*, *Gynecologic Oncology* (1997), thus leaving open the question as to whether the effect of AAV Rep78 was on
15 transcription or translation. The present invention specifically shows that the AAV Rep78 clearly affects p97 transcriptional levels.

The present invention analyzes AAV Rep78 mutants and their ability to bind to DNA and discloses that the favored site for AAV Rep78's binding within the HPV-16 genome is a region within the p97 promoter of the long control region (LCR), from nt 14 to
20 58 using the electrophoretic mobility shift assay (EMSA). These results are surprising as the p97 target sequence contains no GAGC (or GCTC) motifs, which is the core sequence of almost all AAV Rep78 DNA recognitions. This region is important for HPV-16 as it includes functional Sp1 (Gloss, *et al.*, *J. Virol.* 64:5577-5584 (1990), Hoppe-Seyler, *et al.*, *J. Gen. Virol.* 74:281-286 (1993)) and E2 (Androphy, *et al.*, *Nature* 325:70-73 (1987),
25 Moskaluk, *et al.*, *Proc. Natl. Acad. Sci. USA* 85:1826-1830 (1988)) protein binding motifs as well as part of the origin of replication. Furthermore, AAV Rep78 amino acid substitution mutants, at positions 77, 79 or 64-65, identified in this invention, did not recognize p97 DNA. Compared to wild-type AAV Rep78, these AAV Rep78 mutants were found to be defective for inhibition of p97 promoter activity in HeLa and T-47D nuclear
30 extracts *in vitro*, in a transient CAT assay, as well as defective for full inhibition of HPV-16-directed focus formation. These data, taken together, show that the AAV Rep78-p97 promoter interaction is responsible, at least in part, for AAV Rep78-mediated inhibition of

HPV-16. These data indicate that the p97 promoter represents a new prototypical DNA target type for AAV Rep78. Furthermore, AAV Rep78 mutants unable to bind DNA are also defective in their ability to fully inhibit HPV-16 (pL67R)-induced oncogenic transformation, demonstrating the importance of the AAV Rep78-p97 DNA interaction.

- 5 Plasmid L67R contains the oncogene ras coding sequences just downstream of the E7 gene, replacing E1. Thus pL67R is a "super" HPV-16, containing three oncogenes (E6, E7, and ras) expressed from the p97 promoter, instead of just two. Because of its increased activity, the pL67R plasmid allows for easier study of HPV-16 gene expression.

In this patent application, the designations of the AAV Rep78 protein and mutants
10 thereof may also be designated as simply Rep 78 or Rep77^{LG} in the figures and the text.

SUMMARY OF THE INVENTION

The present invention is directed to AAV Rep78 mutants comprising an AAV Rep78 modified protein that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein. Particularly, the AAV Rep78 mutants bind to at least
15 one of PV DNA, AAV DNA, HIV DNA or an oncogene, preferably a promoter sequence, differently as compared to the wild-type AAV Rep78 protein's binding to these DNAs.

The present invention also is directed to fusion proteins containing these AAV Rep78 mutants or wild-type AAV Rep78 proteins, and pharmaceutical compositions containing the AAV Rep78 mutants or fusion proteins comprising the AAV Rep78 mutants or the wild-
20 type AAV Rep78 protein.

The present invention additionally is directed to a method of inhibiting at least one of PV DNA, AAV DNA, HIV DNA or an oncogene, preferably a promoter, whereby PV, PV-associated diseases, cancer, HIV or HIV-associated diseases are inhibited, comprising administering pharmaceutical compositions containing AAV Rep78 mutants or fusion
25 proteins comprising the AAV Rep78 mutants or the wild-type AAV Rep78 protein to patients in need of treatment for these viruses and viral-associated diseases.

The present invention further is directed to a DNA sequence encoding at least one AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein, DNA sequences encoding fusion proteins
30 comprising the AAV Rep78 mutants or the wild-type AAV Rep78 protein and

pharmaceutical compositions comprising a DNA construct containing these DNA sequences encoding the AAV Rep78 mutants or the wild-type AAV Rep78 protein.

The present invention is further directed to a method of treating PV, PV-associated diseases, cancer, HIV or HIV-associated diseases comprising administering the pharmaceutical composition containing a DNA construct containing these DNA sequences encoding the AAV Rep78 mutants that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein or DNA sequences encoding the wild-type AAV Rep78 protein to a patient afflicted with PV-associated diseases, cancer or HIV-associated diseases.

The present invention additionally is directed to a DNA sequence comprising the full length AAV genome (FLAG) modified to replace the AAV Rep78 DNA sequence with the DNA sequence encoding an AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein.

The present invention further is directed to a method of producing rAAV comprising transducing a susceptible mammalian cell with the DNA sequence encoding the minimum AAV genome required to complement a defective rAAV and a DNA sequence encoding the rAAV to obtain increased numbers of rAAV as compared to using AAV sequences containing a wild-type AAV Rep78 genome as a complementor.

The present invention is directed to a method of treating PV-associated diseases, cancer or HIV-associated diseases comprising administering a pharmaceutical composition comprising a AAV Rep78 protein to a patient afflicted with a PV-associated disease or cancer, wherein said AAV Rep78 protein binds to a PV DNA sequence or promoter that controls the expression of a PV oncoprotein or binds to a DNA sequence or promoter that controls the expression of an oncoprotein. Alternatively, the AAV Rep78 protein binds to DNA sequences in the HIV in the HIV genome.

The present invention further is directed to a regulatable element to which AAV Rep78 binds and controls, and a regulatable promoter containing this sequence that preferably includes nucleotides 14-56 as shown in Figure 2

The present invention also is directed to a method of selecting an AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein comprising contacting an AAV Rep78 mutant as described in the present application and contacting the wild-type AAV Rep78 protein with at least one

DNA sequence obtained from one or more of a papillomavirus, an AAV, a HIV or an oncogene for a period of time to allow binding of the mutant and the wild-type Rep78 to the DNA; determining the binding of the mutant and the wild-type Rep78, and compare the binding and select mutants having the desired binding affinities. If numerous binding assays are performed, it may not be necessary to perform the wild-type binding assay each time a Rep78 mutant is tested as the binding of the Rep78 protein will be known.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-D show AAV Rep78 recognition of sequences from the HPV-16 p97 promoter using affinity chromatography and EMSA experiments identifying the region of HPV-16 to which AAV Rep78 binds. The amount of protein added is given in μ g in parentheses. Figure 1A shows the use of AAV Rep78 affinity chromatography to select 32 P-labeled *Pst* I, *Bam* HI double digested HPV-16 DNA. AAV Rep78 preferentially binds a 1.8 Kb *Pst* I fragment from HPV-16 (nt 7003-875). Figure 1B shows that AAV Rep78 preferentially binds sequences from the HPV-16 LCR, nt 7841-106, compared to an equally sized fragment from the MSV-LTR by EMSA analysis. Figure 1C shows that AAV Rep78 preferentially binds nt 14-106 compared to 7841-13 by EMSA analysis. Figure 1D shows that AAV Rep78 preferentially binds nt 14-56 compared to 57-106 by EMSA analysis.

Figure 2 shows the sequences of the HPV-16 p97 promoter with the important elements within the immediate p97 region as labeled boxes.

Figure 3 shows that AAV Rep78 specifically binds the minus (-) strand of p97 using an EMSA analysis of AAV Rep78 interaction with either the + or the - strand of p97 (nt 14-106). The + and - strands were generated by asymmetric PCR amplification.

Figures 4A-B show AAV Rep78 mutant proteins defective in binding AAV TR and the p97 DNA substrates. Figure 4A shows an EMSA analysis of wild-type and mutant AAV Rep78 proteins binding to AAV TR DNA. Note that Rep-64^{LH}65TM does not bind the AAV TR while wild-type and Rep-77^{LG} does. Figure 4B shows an EMSA analysis of wild-type and mutant AAV Rep78 proteins binding to p97 DNA (nt 14-106). Note that in contrast to Figure 4A, both Rep-64^{LH}65TM and Rep-77^{LG} are unable to bind p97.

Figures 5A-B show that AAV Rep78 mutant proteins Rep-64^{LH}65TM and Rep-77^{LG} are defective for inhibiting p97 promoter activity in *in vitro* transcription assays. Shown are representative *in vitro* transcription experiments based on: Figure 5A - HeLa cell nuclear extracts (HPV positive) or Figure 5B - T-47D cell nuclear extracts (HPV negative).

Maltose-binding protein (MBP)-AAV Rep78, Rep-64^{LH65}TM and Rep-77^{LG} were added to the reaction and the generation of the p97 RNA product was determined.

Figure 6 shows that AAV Rep78 mutant genomes FLAG-64^{LH65}TM and FLAG-77^{LG} are defective for inhibiting p97 promoter activity by CAT assay. Cotransfection of dl10-37 (large deletion within AAV Rep78) served as a negative control for inhibition. Cotransfection with increasing doses of pSM620 (2, 4, and 8 μ gs, *wild-type* AAV Rep78) served as a positive control for inhibition. Mutants FLAG-64^{LH65}TM and FLAG-77^{LG} were similarly cotransfected.

Figure 7 shows that AAV Rep78 mutant genomes FLAG-64^{LH65}TM and FLAG-77^{LG} are defective for inhibiting HPV-16 oncogenic transformation. Shown is a representative of three focus formation assays. C127 contact inhibited murine fibroblasts were calcium phosphate transfected with 3 μ gs of pL67R plus 6 μ gs of the indicated AAV plasmid. dl10-37 and pSM620 are the controls, encoding a large deleted AAV Rep78 and wild-type AAV Rep78 respectively. Note that FLAG-64^{LH65}TM and FLAG-77^{LG} are significantly defective in inhibiting pL67R oncogenic transformation compared to pSM620.

Figures 8A-B show that AAV Rep78-p97 interaction is not as strong as AAV Rep78-TR interaction. Figure 8A shows a competitive EMSA experiment analyzing wild-type and 77^{LG} AAV Rep78 proteins binding to a ³²P-labeled TR substrate with competition from unlabeled TR and p97 DNA. Three doses of competitor DNA (0.1, 0.5, and 1 μ g) were added as indicated by the triangle. Note that TR DNA is a better competitor than p97. Figure 8B shows a competitive EMSA experiment of wild-type AAV Rep78 protein binding to a ³²P-labeled p97 substrate with competition from unlabeled TR and p97 DNA. Three doses of competitor DNA (0.1, 0.5 and 1 μ g) were added as indicated by the triangle. Again, note that TR DNA is a better competitor than p97.

Figures 9A-B show that AAV Rep78 binds sequences of the BPV-1 long control region (LCR). Figure 9A shows an EMSA in which AAV Rep78 binds to the BPV-1 LCR (nt 7758-7030). Note that the DNA-protein complex occurs in a dosage dependent manner with increasing addition of MBP-AAV Rep78. These LCR sequences contain two E2 motifs. Figure 9B shows a competitive EMSA demonstrating that the AAV TR is a better competitor than the BPV-1 LCR itself. 0.1 μ g of MBP-Rep were added as indicated. Four doses of synthetic competitor DNA (Durst, *et al.*, *Proc. Natl. Acad. Sci. USA* 80:3812-

3815 (1983), 5, 20 and 40 ng) were added as indicated by the triangle. Note that the MSV-LTR DNA is not an effective competitor compared to the BPV LCR DNA.

Figure 10 shows the *in vitro* transcription showing AAV Rep78 mutant inhibition of transcription.

5 Figure 11 shows that FLAG-77^{LG} replicates at higher levels and that FLAG-192^{HG} replicates at slightly depressed levels compared to wild-type.

Figure 12 shows that Rep-192^{HG} binds AAV TR (terminal repeat) DNA in a different higher order complex and with higher affinity.

10 Figure 13 shows that Rep-192^{HG} binds to the p5 promoter sequences in a higher order complex than wild-type Rep78.

Figure 14 shows the effect of several AAV Rep78 mutants on the inhibition of HPV-16-induced oncogenic foci.

Figure 15A-C discloses the nucleotide sequence of the AAV2 genome as disclosed in Srivastava, A. *et al.*, *J. Virol.* 45:555 (1983). The nucleotide sequence encoding AAV
15 Rep78 is nucleotides 321-2186, in which nucleotides 2184-2186 are the stop codon, taa.

Figure 16 discloses the 621 amino acid sequence of AAV Rep78 encoded by nucleotides 321-2183 as discussed above in the description of Figure 15.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

20 AAV Rep78 is responsible for all enzymatic functions of the AAV life cycle as well as for the inhibition of HPVs. The present invention is directed to AAV Rep78 mutants comprising a modified AAV Rep78 protein that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein. These different functions can be enhanced or defective biochemical or biological functions as compared to
25 the wild-type AAV rep78 protein. The mutants of the present invention also are useful to map the function domains of Rep 78.

More specifically, the modified AAV Rep78 protein or AAV Rep78 mutants bind to at least one DNA sequence obtained from a PV, an oncogene, an AAV an HIV or a combinations thereof differently than the wild-type AAV Rep78 protein binds to these
30 DNAs. The mutants are useful for the inhibition of viruses, particularly, PVs and HIV, and therefore, these mutants will be useful in the treatment of PV-associated diseases, preferably those disease caused by a HPV or BPV, but also other PVs known to persons

skilled in the art are intended to be encompassed by the present invention. The mutants also are useful for treating the diseases that result from the expression of an oncogene or HIV DNA.

The present invention is intended to encompass two different functional types of AAV Rep78 mutants. The first type of AAV Rep78 mutant has enhanced strong binding to PV DNA or oncogenes or AAV DNA or HIV DNA or a combination of one or more of these DNAs, preferably promoter regions of these DNAs, and provides enhanced control of PV, oncogene and HIV expression. The second type of AAV Rep78 mutant has weak or no binding affinity to PV or AAV DNA or both, yet has other intact AAV functions and provides for improved AAV and rAAV production, and controls PV by increasing the numbers of AAVs.

The first type of AAV Rep78 mutant, typified by Rep192^{HG} and phenotypically related Rep proteins that bind similarly to Rep192^{HG}, possesses enhanced DNA binding to at least one of a PV DNA, an AAV DNA or HIV DNA or an oncogene as compared to the wild-type AAV Rep78 protein's binding to these DNAs. This latter mutant type inhibits PV or other target genes by inhibiting the transcriptional activity or promoter of this virus or other target genes, and also inhibits the transcriptional activity or promoters that drive the expression of oncoproteins. Enhanced promoter binding directly correlates with enhanced promoter inhibition (See Figures 10, 14). Therefore, these latter mutants are useful in anti-PV and anti-cancer protocols, not only for treating PV, but also for inhibiting the oncogenic transformation of the cell in general. This mutant type is demonstrated by the Rep-192^{HG} and is disclosed in the present invention. But other similar mutants may be selected using the inhibition assays described in the present invention.

The second type of AAV Rep78 mutant is a mutant possessing weak or no DNA binding affinity to at least one DNA sequence obtained from a PV or an AAV, such as the promoter, AAV p5 or HPV-16 p97, when this binding affinity is compared to the binding of the wild-type AAV Rep78 protein. Although these mutants bind less strongly, if at all, to the selected DNA(s), these mutants have other functions that are intact, and therefore have enhanced ability to complement AAV functions, that results in the generation of higher levels of AAV DNA replication and/or AAV virion numbers. These increased levels of AAV DNA replication and virions are useful for generating more rAAV for gene therapy as compared to the wild-type Rep78. But these weak or no binding AAV Rep78

mutants, also are useful in treating PV by virtue of the negative impact that the presence of AAV has on the presence of PV. Thus, any mechanisms that would increase AAV numbers is expected to decrease PV. The Rep-77^{LG}, Rep-79^{FA} and the Rep-64^{LH}65TM mutants are examples of this type of mutant and are disclosed in the present invention.

5 In another embodiment, the present invention is directed to AAV Rep78 mutants that include 1) truncated wild-type AAV Rep78 proteins, 2) wild-type AAV Rep78 proteins containing amino acid substitutions, 3) wild-type AAV Rep78 proteins containing internal amino acid deletions, and 4) wild-type AAV Rep78 proteins that contain combinations of these modifications. The AAV Rep78 mutant in one embodiment is a modified AAV
10 Rep78 protein containing at least the minimum number of amino acids of the wild-type AAV Rep78 protein necessary to bind to at least one of the PV DNA, an oncogene or the AAV DNA, or HIV DNA, preferably the promoter region of these DNAs, to obtain enhanced inhibition of PV or oncoproteins or HIV. In the case of the oncogene encoding the oncoprotein, the proto-oncogene promoter is inhibited. In certain mutants, this
15 modification could reduce the binding affinity. If enhanced binding is the property of the AAV Rep78 mutant, then the goal of the present invention is to select the smallest truncated AAV Rep78 mutants that are still capable of binding to the PV DNA, oncogene, AAV DNA, HIV DNA or a combination thereof, to inhibit the expression of PV, PV oncoproteins or other cancer related oncoproteins or HIV to inhibit transcription with the
20 end result being the inhibition of the PV-associated disease or cancer or HIV-associated disease. Again, if enhanced binding to the promoter region is the goal, then the truncated AAV Rep78 mutant must still maintain the ability to bind to the promoter region of at least one of PV DNA or AAV DNA or oncogene or HIV DNA, and particularly bind to nucleotides 14-56 of the p97 of HPV-16. Truncated Rep 78 mutants, within the context of
25 the present invention, include any deletion of amino acids of the wild-type Rep78 protein. These deletions may be serial deletions of amino acids from the N-terminus, middle or C-terminus of the protein or one or more amino acid deletions from different portions of the protein. A further embodiment is directed to a truncated AAV Rep78 mutant that is composed of at least two of these truncated wild-type AAV Rep78 linked to each other to
30 form a multimer AAV Rep78 mutant. Protein fusions are performed by techniques known in the field or are made by recombinant techniques by linking the DNAs encoding the proteins with the resulting expression of a fusion protein.

The present invention provides sufficient guidance to a person skilled in the art to obtain additional AAV Rep78 mutants that have modified binding affinities as compared to the wild-type AAV Rep78. The nucleic acid sequence is disclosed in Srivastava *et al.* *J. Virol.* 45:555 (1983) and is disclosed in Figure 15A-C which is used to prepare mutants according to the present invention. AAV has been studied and the organization of the AAV genome is disclosed in Muzyczka, N. *et al.*, *Current Topics in Microbiology and Immunology* 158: 97-129 (1992), particularly Figure 3. The present invention discloses a sequence to which the AAV Rep78 binds to in Figure 2 and discloses methods to test for the strength of the binding or the lack of binding to DNA sequences, preferably promoter regions. The present invention is intended to encompass other AAV Rep78 mutants that have modified DNA binding; for example, promoter binding affinities. Such methods would not require undue experimentation and are disclosed in the present invention or known to skilled persons.

In a further embodiment, the present invention is directed to a fusion protein comprising an AAV Rep78 mutant described above. The fusion protein envisioned in the present invention is one in which the *tat* protein of HIV is linked to the AAV Rep78 mutant or the wild-type AAV Rep78. Particularly, the *tat* protein of HIV type 1 is preferred for linking to the AAV Rep78 wild-type protein or mutants to provide a fusion protein for direct treatment of the Rep78 mutant and the *tat* protein of HIV to cells and tissues. The method circumvents the need for introducing the gene encoding the AAV Rep78 mutant via gene therapy. The *tat*-wild-type Rep78 or *tat*-Rep78 mutant, optionally as a fusion protein with MBP (*E. coli* maltose binding protein), is able to be actively taken up by cells. This method provides for these proteins containing *tat* to be taken up and allows AAV Rep78 and its mutants to be applied by cream or other forms of an external pharmaceutical compositions for dermatological application. Further, these fusion proteins also are applied by direct injection into the tumor sites for a potentially higher effectiveness or administered directly to the tumor if visible from the surface of the skin. Nagahara *et al.*, *Nature Medicine*, 4(12):1449 1998 discloses the use of *tat* in fusion proteins.

The present invention is also directed to a fusion protein comprising a maltose-binding protein (MBP) linked to said AAV Rep78 mutant. See Batchu *et al.*, *Biochem. Biophys. Res. Commun.* 208: 714 (1995).

The present invention further is directed to pharmaceutical compositions comprising at least one AAV Rep78 mutant or AAV Rep78 protein as disclosed above in admixture with a pharmaceutically acceptable carrier. The preferred manner of treatment using these pharmaceutical compositions is subcutaneous injection or directly into the tumor or via
5 topological administration using creams or lotions that may include lipofection where the mutant is carried in beads made of bipolar lipids. Pharmaceutical compositions and appropriate carriers for each of these modes of administration are known to persons skilled in the art and can be combined without undue experimentation with the AAV Rep78 proteins and mutants of the present invention. See, for example, Remington's Pharmaceutical Sciences:
10 Drug Receptors And Receptor Theory, (18th ed.), Mack Publishing Co., Easton, PA (1990) These pharmaceutical compositions are administered to inhibit PV-associated diseases, cancer and HIV-associated diseases in patients having these diseases.

AAV as well as adenoviruses and retroviruses, are useful as gene transfer vehicles in gene therapy. Gene therapy techniques have been known for several years, and as
15 discussed in Kay *et al.*, *Proc. Natl. Acad. Sci. USA*, 94:12744 (1997), vehicles for gene transfer can be classified into two major classes: viral and nonviral vectors. Table 1 of Kay *et al.* discloses retroviruses, adenoviruses and adeno-associated viruses (AAV) as viral vehicles and liposomes as nonviral vehicles. The drawbacks of using retroviruses and adenoviruses are that they are large and as such, carry extra genes into the transduced cells,
20 which encode additional proteins that likely will elicit an immune response from the patient. AAV is smaller in size and provides less complicated epitopes to the patient than retroviruses or adenoviruses. Both AAV and retroviruses have been demonstrated as being highly effective gene transmitters in immortalized cells. However, retroviruses are unable to transduce non-dividing cells. Furthermore, there is another very important difference
25 between the two viruses in regard to their use as a human gene therapy vector. AAV is non-pathogenic. In contrast, retroviruses are pathogenic. This fact was recently emphasized in monkey bone marrow gene therapy experiments utilizing retrovirus based vectors, even minute amounts (10^{-6} of the vector) of contaminating wild-type retrovirus resulted in the most of the animals developing malignant lymphoma (Donahue *et al.*, *J. Experimental Medicine* 176(4): 1125 (1992); Vanin *et al.*, *J. Virol.* 68(7): 4241 (1994)).
30

AAV actually has anti-cancer properties. Therefore, AAV is a suitable choice for transducing papilloma tumor cells, such as cervical cancer or other cancers with which HPV is associated (Hermonat *et al.*, *Proc. Natl. Acad. Sci USA*, 81: 6466 (1984)).

In a further embodiment, the present invention is directed to a DNA sequence
5 encoding at least one AAV Rep78 mutant or AAV Rep78 protein described in this application. Additionally, the DNA sequence encoding an AAV Rep78 protein or a mutant thereof is further contained in a vector for delivery of the virus to the target cell in culture or in a patient's body that will benefit from treatment of AAV Rep78 or a mutant thereof. This vector is any known vector and for gene therapy is preferably an AAV vector but
10 other known vectors, such as retroviruses, adenoviruses, pox viruses or liposomes are also useful in the present invention. This vector containing the DNA sequence is admixed with a pharmaceutically acceptable carrier for administration to a patient afflicted with a PV, PV-associated disease, cancer, HIV or HIV-associated disease. This method of treatment delivers DNA encoding AAV Rep78 protein or an AAV Rep78 mutant to the PV tumor of
15 the patient so that AAV Rep78 DNA or its mutant DNA transduces the cells and is expressed by the affected cells causing an inhibition of PV production or oncoproteins. The vector is useful for gene transfer: *ex vivo* where cells are removed, genetically modified by transduction with the vector, and transplanted back into the same recipient or *in vivo* therapy accomplished by transfer of genetic materials directly into the patient's cells
20 in the body. Methods of constructing and using such vectors are known in the art. Hermonat, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 81:6466 (1984); McLaughlin *et al.*, *J. Virology* 62(6):1963 (1988); Hermonat *et al.*, *FEBS Letters* 407:78 (1997); Liu *et al.*, *J. Inf. Cytok. Res.* 20:21-30. (2000); Hermonat *et al.*, *J. Hum. Virol.* 3: 113-124 (2000).

The present invention is further directed to a DNA construct comprising a DNA
25 sequence encoding AAV Rep78 or a AAV Rep78 mutant operably linked to an inducible promoter, preferably a promoter that is induced by a molecule produced during PV infection or cancer, such as an oncoprotein. As discussed above, this DNA construct is useful for insertion into a DNA vector, such as an AAV vector, retrovirus, or other vectors as discussed above, for transfer into mammalian cells, when admixed with a
30 pharmaceutically acceptable carrier for treating a patient and inhibiting PV-associated disease in the patient. Examples of such inducible promoters that are functional in the present invention are the BPV p89 promoter that is responsive to the PV E2 protein and the

promoter of the E2 gene of adenovirus which is responsive to the HPV E7 oncoprotein. Other appropriate promoters inducible by specific PV proteins may be selected by persons skilled in the art.

The present invention also is directed to a DNA sequence comprising the full length
5 AAV genome (FLAG) modified in that the AAV Rep78 gene is replaced with an AAV Rep78 mutant of the present invention.

The present invention further is directed to a DNA sequence comprising at least the minimum portion of the AAV genome sufficient to complement a defective rAAV, wherein the portion of the AAV genome is modified to replace the AAV Rep78 DNA sequence with
10 the DNA sequence encoding the AAV Rep78 mutants described herein, such as the AAV Rep-77^{LG} and Rep-79^{FA} mutants, which possesses no DNA binding or weak DNA binding to the DNA sequence obtained from at least one of a PV or an AAV or an oncogene, preferably the promoter regions of these sources, that results in the generation of higher levels of rAAV DNA replication and virion numbers. This DNA sequence is useful in
15 producing rAAV comprising transducing a susceptible mammalian cell with the minimum portion of the AAV genome sufficient to complement a defective rAAV as a complementor, and a DNA sequence encoding the rAAV to obtain increase numbers of rAAV or AAV. The rAAV is the AAV with all of the AAV genes removed except the TR sequences (terminal repeats), which renders the rAAV defective, and in need of a helper virus as a
20 complementor, to provide the missing AAV genes for replication. The rAAV also contains a DNA sequence encoding a heterologous gene. The DNA sequence sufficient to complement a defective rAAV comprises at least the AAV *lip-cap* gene and the DNA sequence encoding an AAV Rep78 mutant that binds weakly or not at all to DNA sequences, preferably a promoter region from at least one PV, an oncogene, AAV or HIV.

25 The present invention further is directed to a method of inhibiting PV, PV-associated diseases, cancer, HIV or HIV-associated diseases comprising administering a pharmaceutical composition comprising an AAV Rep78 protein or a mutant thereof, to a patient afflicted with a PV-associated disease, wherein the AAV Rep78 protein or mutant binds to HPV DNA, particularly binds to nucleotides 14-56 of p97 of HPV-16, thereby
30 inhibiting expression of HPV oncoproteins and/or binds to HIV DNA, for example, such as HIV promoter sequences TAR or nucleotides -54 to -34.

The present invention also discloses an AAV Rep78 regulation element comprising the nucleotides shown in the nucleotide sequence of Figure 2, wherein this element provides the binding site for the AAV Rep78 protein, and particularly comprises about nucleotides 14 -56 of the nucleotide sequence of Figure 2. This present invention also discloses an AAV Rep78 regulatable promoter comprising the regulation element of the nucleotides shown in the nucleotide sequence of Figure 2, wherein this element provides the binding site for the AAV Rep78 protein, and particularly comprises about nucleotides 14 -56 of the nucleotide sequence of Figure 2. This AAV Rep78 regulatable promoter comprises the regulation element described above and the remaining promoter sequences from a promoter other than the HPV-16 p97 promoter.

The following specific examples provide guidance to the skilled person regarding the methods useful in preparing and testing AAV Rep78 mutants for their capability and relative affinity in binding PV DNA and AAV DNA and testing the effects of that binding or lack thereof on transcription. These examples should by no means be interpreted as limiting the present invention to these disclosed examples but as a guide to select additional AAV Rep78 mutants that fall within the spirit of the present invention.

EXAMPLES

The following experimental procedures are utilized in the present invention:

Experimental Procedures

20 *AAV Rep78 affinity chromatography selection of HPV-16 DNA fragments*

The *Bam* HI fragment containing the complete HPV-16 genome (without plasmid sequences, from pAT/HPV-16) was isolated by gel electrophoresis using the GeneClean II kit. This DNA was further digested with *Pst* I, Klenow labeled with $\alpha^{32}\text{P}$ -dCTP, and cleaned by phenol extraction and ethanol precipitation. The AAV Rep78 affinity chromatography was carried using 50 μg s of MBP-AAV Rep78 bound to 100 μl s of amylose resin (New England Biolabs). Klenow labeled ^{32}P -HPV-16 DNA (2 μg s), digested by *Pst* I and *Bam* HI, were applied to the column in 100 μl s of column buffer (20mM Tris [pH 7.4], 200 mM NaCl, 1 mM EDTA, 1 mM DTT) and incubated for 15 minutes at room temperature. After washing the column twice with 1 ml of column buffer, the bound DNA was eluted with 100 μl s of 1% SDS, 20 mM Tris pH7.5. The eluted products were then analyzed by PAGE (4%), dried and autoradiographed.

DNA substrates and the electrophoretic mobility shift assay (EMSA)

HPV-16 and MSV-LTR DNA substrates were generated by polymerase chain reaction (PCR) amplification. The AAV terminal repeat (TR) substrate was generated by the ligation of three separate synthetic oligonucleotides as described previously (Bishop, *et al.*, *FEBS Letters* 397:97-100 (1996)). The TR was 5' end labeled with polynucleotide kinase using ³²P ATP (5,000Ci/mmol, Amersham). Single stranded DNA substrates were generated by asymmetric PCR amplification as previously described (Gyllenstein, *et al.*, *Proc. Natl. Acad. Sci. USA* 85:7652-7656 (1988)). EMSA was carried out as follows; approximately 1ng of ³²P-labeled DNA substrate was incubated with increasing amounts of MBP-AAV Rep78 for 10 min. at room temperature in binding buffer [(25 mM HEPES KOH pH 7.5, 10 mM MgCl₂, 1mM dithiothreitol, 2% glycerol, 25 ug bovine serum albumin, 50 mM NaCl, 0.01 % NP40 and 0.5 ug poly (dI-dC)]. Samples were electrophoresed in a 4% polyacrylamide gel (40:1 acrylamide and bis-acrylamide weight ratio) with 5% glycerol in 0.5x TBE buffer at 100V for about 3 hr. Gels were dried and autoradiographed at -70°C.

Construction of AAV Rep78 mutant plasmids and production of MBP-AAV Rep78

chimeric proteins The construction of any of the AAV Rep78 mutants utilizes the known AAV Rep78 nucleic acid sequence as disclosed, for example, in Figure 15A-C and in Srivastava *et al.*, *J. Virol.* 45:555 (1983). Further, the 621 amino acid wild-type Rep78 encoded by nucleotides 321-2183 (stop codon nucleotides 2184-2186) is disclosed in Figure 16 and was known. The construction of the plasmid pMAL-Rep-64^{LH65}TM from which mutant MBP-64^{LH65}TM protein is produced, has been described previously (Batchu, *et al.*, *Biochem. Biophys. Res. Comm.* 208:714-720 (1995), Batchu, *et al.*, *Biochem. Biophys. Res. Comm.* 210:717-725 (1995)). The plasmid pMal-Rep-77^{LG} was similarly constructed using a different mutagenic oligonucleotide (5'-CCCCGGAGGCCGAATTCTTTGTGCAA) and the M13 based plasmid pALTER-AAV3 (containing all of the AAV genes). A second oligonucleotide created a *Sph* I restriction site immediate upstream of AAV Rep78 ORF. The mutations were initially characterized by the generation of a new restriction site and were further verified by DNA sequencing with Sequenase (U.S. Biochemicals) according to the manufacturer's recommendations. The mutant AAV Rep78 ORFs were then transferred into pMALc2 on an *Sph* I and *Xho* I fragment (nt. 321 to nt. 2233) to generate pMAL-Rep-64^{LH65}TM and pMAL-Rep-77^{LG}. Both the mutant and wild-type fusion proteins with MBP

were purified by affinity chromatography using amylose resin following kit directions (Protein Purification and Expression System, New England Biolabs). Fractions were collected and analyzed by SDS-polyacrylamide gels. Purified fractions were concentrated using Centricon 10 kDa cut-off membrane filters (Amicon). Routinely these procedures
5 resulted in MBP-AAV Rep78 and 64^{LH65}TM proteins of 70-90% purity with a yield of 20 µg/100ml bacterial culture.

Construction of full length AAV genomes (FLAG) carrying the AAV Rep78 mutations

The *Bsa* I fragment (4.2 Kb, containing all of the AAV genes) from pALTER-AAV-64^{LH65}TM and pALTER-AAV-77^{LG} were isolated and ligated to the *Bsa* I partial digestion
10 fragment (4.9 Kb, containing pBR322 plus the AAV TRs) from pSM620 (Samulski, *et al.*, *Proc. Natl. Acad. Scii. USA* 79:2077-2081 (1982)) to generate pFLAG-64^{LH65}TM and pFLAG-77^{LG}. To insure that the mutations were transferred into the full length AAV background, the region of the mutation was once again sequenced as described above.

In vitro transcription analysis of p97 promoter activity

15 An HPV-16 p97-CAT DNA fragment was used as a template for transcription. The p97-CAT DNA fragment was generated by PCR amplification using primer 1 (5'ACAAGCAGGATTGAAGGCCA, HPV-16 nt 7043-7065) complementary to the p97 sequences) and primer 2 (5' CATATCACCAGC TCACCGTC, nt 615-633 of pSV2CAT) complementary to the CAT sequences). The plasmid p16P (p97-CAT) was used as the
20 original PCR template (Romanczuk, *et al.*, *J. Virol.* 64:2849-2859 (1990)). This produced a 1.2 Kb product. A 25 µl reaction mixture contained 0.5 µg of DNA template, 20 mM HEPES pH 7.9, 5 mM MgCl₂, 100 mM KCl, 0.5 mM DTT, 20% glycerol, 25 µM [³²P] GTP, 400 µM ATP, CTP and UTP, and 8 units Hela nuclear extract (Promega, HPV positive cervical cancer) or 5 µg of T-47D nuclear extract (Geneka Corp., HPV negative
25 breast ductal carcinoma). Reactions were incubated at 30°C for 60 min, and then terminated by adding 175 µL of Stop Solution containing 300 mM Tris-HCl, pH 7.9, 0.5% SDS, 300 mM sodium acetate, 2 mM EDTA and 3 µg/ml tRNA. RNA was extracted with phenol-chloroform, precipitated with ethanol, and finally dissolved in 10 µL of formamide containing 0.1% each of xylene cyanol and bromophenol blue. Samples were analyzed on
30 an 6% polyacrylamide, 7 M urea gel, dried, analyzed by autoradiography. An p97-specific RNA product of approximately 300 bases results.

Transient chloramphenicol acetyltransferase assay

Transient CAT assays were carried out by calcium phosphate transfection of the p16P (p97-CAT) plasmid plus several AAV plasmids (amounts indicated within the Figure
 5 legenda). Forty eight hours after transfection cell extracts were prepared, equalized for protein content by spectroscopic analysis at 280 nm, and assayed as described previously (Romanczuk, *et al.*, *J. Virol.* 64:2849-2859 (1990), Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994)).

Focus formation assay

Contact inhibited murine C127 mouse fibroblasts were calcium phosphate
 10 transfected with 3 µgs of the HPV-16/*ras* chimeric plasmid, pL67R (Schmitt, *et al.*, *Virology* 172:73-81 (1989)) plus 6 µgs of either pSM620 (+AAV Rep78, *wild-type*), d110-37 (-AAV Rep78, large deletion)(Hermonat, *et al.*, *J. Virol.* 51:329-333 (1984)), pFLAG-64^{LH}65TM, or pFLAG-77^{LG}. The plasmid pL67R contains the EJ-H-*ras* coding sequences ligated in place of the E1 gene (Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994)).
 15 Thus, pL67R contains three oncogenes (E6, E7, and *ras*), all of which are expressed from the p97 promoter. The cells were fed for two and one half weeks, fixed with formaldehyde and stained with methylene blue.

The following experiments were performed using the procedures described above:

AAV Rep78 recognizes the p97 promoter DNA

20 To observe if AAV Rep78 bound to HPV-16 DNA, AAV Rep78 affinity chromatography was utilized. MBP-AAV Rep78, bound to amylose resin, was incubated with ³²P-labeled HPV-16 DNA fragments generated from *Pst*I and *Bam* HI double digestion. After washing and elution the products were agarose gel electrophoresed adjacent to unselected HPV-16 fragments. As shown in Figure 1A, although a few partial
 25 digestion products are present, it was clear that it was the 1.8 kb *Pst*I fragment (nt 7003-875) of HPV-16 which was preferentially recognized by AAV Rep78.

Within the 1.8 kb fragment lies the long control region (LCR) of HPV-16 which contains central *cis* elements (origin of replication [*ori*], enhancers and promoters) essential for HPV-16 biological function. Furthermore, AAV Rep78 is a viral transcription factor
 30 known to bind promoter DNA (McCarty, *et al.*, *J. Virol.* 74:4988-4997 (1994), Batchu, *et al.*, *Cancer Letters* 86:23-31 (1994), Wonderling, *et al.*, *J. Virol.* 71:2528-2534 (1996)). Thus, it is reasoned that AAV Rep78 might be targeting the *ori*/p97 region within this

fragment due to AAV Rep78's known modulation of the HPV-16 p97 and HPV-18 p105 promoters (Hermonat, P.L., *Virology* 172:253-261 (1989), Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994), Hermonat, *et al.*, *Gynecologic Oncology* 66:487-494 (1997), Horer, *et al.*, *J. Virol.* 69:5485-5496 (1995)), and of BPV-1 DNA replication (Hermonat, P.L. *Virology* 189:329-333 (1992)). To map the region of binding sequentially smaller substrates from this region were tested for recognition by AAV Rep78 (Figures 1B-C) by EMSA analysis. In Figure 1B AAV Rep78 was shown to strongly recognize the HPV-16 sequences from nt 7814-106 (p97), while it does not significantly recognize a similarly sized analogous fragment from the MSV-LTR. These data clearly indicate that there is a specific recognition of the p97 DNA well above non-specific binding. As mentioned earlier, AAV Rep78 is able to inhibit expression from p97, but has little effect on the MSV-LTR (Hermonat, P.L., *Cancer Research* 51:3373-3377 (1991)). Thus, AAV Rep78 binding of promoter DNA may be associated with an ability to regulate p97 expression. In Figure 1C, the target sequence was further defined to be in the 3' half of this region (nt 14-106, hereafter referred to as "p97"). Finally, in Figure 1D, a strong target sequence for AAV Rep78 binding is shown to be contained within nt 14-56. Figure 2 shows the sequences of this region. Note that the nt 14-56 sequences contain an intact E2 binding motif and an Sp1 binding motif. These sequences also partially overlap the HPV-16 *ori* and E1 binding regions.

The E2 motifs are interrupted palindromes and can potentially form hairpin structures. There are also other interrupted palindromic sequences present within the p97 promoter region (nt 14-106). The DNA substrate which has formed some degree of secondary structure (2ndS) was observed in the EMSA gels as an elevated band over the lower simple duplex (dup) DNA substrate. It is this higher band which is the preferred substrate for AAV Rep78 recognition as demonstrated most clearly in Figure 1B. This preference for DNA substrates with secondary structure has also been seen in AAV Rep78 recognition of the TAR region DNA of HIV type 1 (Batchu, *et al.*, *FEBS Letters* 367:267-271 (1995)). In order to test the possibility that AAV Rep78 was recognizing secondary structure, the + and - strands of the p97 sequence were separately generated by single sided PCR. Such single stranded DNA should naturally form secondary structure as single stranded RNA does. The ³²P-labeled + and - strands were compared by EMSA for AAV Rep78 recognition as shown in Figure 3. As can be seen the - strand was strongly

recognized by AAV Rep78, while the + strand was not. These data suggest that AAV Rep78 may be recognizing both secondary structure and the specific sequence of the DNA. *Two AAV Rep78 amino acid substitution mutants are defective for binding p97 promoter DNA* AAV Rep78 mutant proteins with specific amino acid substitutions for study in

5 dissecting AAV Rep78's functions and domains are generated. The mutant AAV Rep78 proteins were produced as fusions with the maltose binding protein (MBP) as previously described (Batchu, *et al.*, *Biochem. Biophys. Res. Comm.* 208:714-720 (1995), Batchu, *et al.*, *Biochem. Biophys. Res. Comm.* 210:717-725 (1995)). During the characterization of one MBP-AAV Rep78 mutant protein, Rep-77^{LG} (substituting a glutamine for a leucine at

10 amino acid 77), it was found to be able to bind an AAV terminal repeat (TR) DNA substrate at levels comparable to wild-type MBP-AAV Rep78, as shown in Figure 4A. In contrast Rep-77^{LG} was defective in recognizing the p97 promoter as shown in Figure 4B. In these experiments the wild-type MBP-AAV Rep78 and Rep-64^{LH65}TM proteins served as the positive and negative controls, with Rep-64^{LH65}TM being unable to bind any DNA

15 substrate thus far assayed by EMSA (Batchu, *et al.*, *Biochem. Biophys. Res. Comm.* 210:717-725 (1995), Kokorina, *et al.*, *J. Hum. Virol* 1:441-450 (1998)). Thus, both Rep-64^{LH65}TM and Rep-77^{LG} were defective for binding p97. However, Rep-77^{LG} was particularly interesting as it was able to distinguish between the TR and p97 substrates. Thus, the mechanism of recognition used by AAV Rep78 to bind the AAV TR was

20 different, at least in part, than that for recognizing p97.

AAV Rep78 mutants defective for binding p97 are also defective for inhibition of p97 promoter activity The effects of AAV Rep78 protein DNA binding on p97 promoter activity was observed. *In vitro* transcription was used to study these protein affects. Thus, various amounts of all three proteins (*wt*, Rep-64^{LH65}TM, and Rep-77^{LG}) were added to

25 HeLa and T-47D cell nuclear extracts and containing a p97 DNA template. The experiments were done four times using HeLa extracts and twice using T-47D extracts, all giving similar results. Representative experiments are shown in Figure 5. As seen, the addition of increasing amounts of maltose-binding protein (MBP)-AAV Rep78 inhibited RNA initiation from the p97 promoter in a dosage dependent manner in both the HeLa and

30 T-47D extracts. In sharp contrast to MBP-AAV Rep78, the addition of increasing amounts of the AAV Rep78 mutants, AAV Rep-64^{LH65}TM and AAV Rep-77^{LG}, protein had little or

no effect on p97 promoter activity in both cell types. Thus, both AAV Rep78 mutants defective for binding p97 DNA were also defective for inhibition of p97 promoter activity.

A CAT assay was also utilized to further verify AAV Rep78's effects upon the p97 promoter and the results are shown in Figure 6. SW13 breast cancer cells were calcium phosphate transfected with the 4 µgs of plasmid p16P which contains the CAT coding sequences expressed from the HPV-16 p97 (Romanczuk, *et al.*, *J. Virol.* 64:2849-2859 (1990)) plus different amounts of one of four AAV plasmids. FLAG plasmids, pSM620 (wild-type AAV Rep78), FLAG-64^{LH}65TM, and FLAG-77^{LG} were co-transfected with p16P. As indicated by their names the FLAG plasmids contained the same mutated AAV Rep78 sequences as were present in the pMal-based plasmids used to generate the mutant proteins. The basal expression control plate was transfected with the AAV plasmid dl10-37 which contains a large deletion within the AAV Rep78 sequences and served as a negative control. Cotransfection with increasing doses of pSM620 (2, 4, and 8 µgs, wild-type AAV Rep78) served as a positive control for inhibition. Mutant FLAG-64^{LH}65TM and FLAG-77^{LG} were similarly cotransfected. Two days after transfection cellular extracts, equalized for protein content, were assayed for CAT activity. Note that pSM620 was able to inhibit p97 activity, while FLAG-64^{LH}65TM and FLAG-77^{LG} were defective. The results shown in Figure 6 demonstrate, similar to the *in vitro* transcription assays, that AAV Rep78 (pSM620) inhibits p97 activity while FLAG-64^{LH}65TM and FLAG-77^{LG} were defective and did not inhibit p97 activity.

AAV Rep78 mutants defective for binding p97 are also defective for full inhibition of HPV-16-induced oncogenic transformation To further observe the biological significance of AAV Rep78-p97 interaction, the AAV Rep78 mutants (Rep-77^{LG} and Rep-64^{LH}65TM), which were unable to bind p97 and unable to inhibit p97 transcription, were analyzed for their ability to inhibit HPV-16 p97-directed oncogenic transformation. The FLAG-77^{LG} and FLAG-64^{LH}65TM mutant genomes were then tested in a C127 cell-based, HPV-16-induced focus formation assay. An HPV-16/*ras* chimeric plasmid, pL67R, in which the E1 gene is replaced by the EJ-H-*ras* coding sequences was utilized (Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994), Hermonat, *et al.*, *Gynecologic Oncology* 66:487-494 (1997)). This construct, expressing three oncoproteins from the p97 promoter, has higher transforming activity than HPV-16 alone (Hermonat, P.L., *Cancer Research* 54:2278-2281 (1994)). dl10-37 (AAV Rep78 negative) and pSM620 served as the negative

and positive controls, respectively. Plates of C127 cells were calcium phosphate transfected with pL67R (3 µgs) plus one of the indicated AAV plasmids (6 µgs). After 2.5 weeks the cells were formaldehyde fixed, methylene blue stained, and the foci counted. The results are shown in Figure 7. Note that both FLAG-64^{LH}65TM and FLAG-77^{LG} were defective when compared to pSM620 (*wild-type* AAV Rep78) for inhibiting pL67R. Two additional transfection sets gave similar results. As transformation by L67R is dependent on the p97 promoter, these data strongly suggest that the AAV Rep78-p97 interaction is important for AAV Rep78's inhibitory ability.

AAV Rep78-p97 interaction is not as strong as AAV Rep78-TR interaction

To compare the affinities of AAV Rep78 for the AAV TR and HPV-16 p97 DNA substrates a series of competitive EMSA experiments were undertaken. As shown in Figure 8A unlabeled TR DNA competitor was able to strongly inhibit *wild-type* AAV Rep78-³²P-TR interaction, while unlabeled p97 DNA (nt 14-106) competitor was not. Unlabeled p97 DNA competitor was also not able to inhibit Rep-77^{LG}-TR interaction. In Figure 8B it is shown that both unlabeled TR and p97 competitors are able to successfully inhibit *wild-type* AAV Rep78-³²P-p97 interaction, but that the TR is the more effective competitor. These data are consistent with AAV Rep78 having a higher affinity for the AAV TR (its natural substrate with GCTC³) than the HPV-16 p97 promoter (lacking GCTC motifs). These data are also consistent with the interpretation of the other data, Figure 5, that the mechanism of AAV Rep78 recognition of p97 DNA is different than AAV Rep78 recognition of TR DNA.

The BPV-1 LCR also contains multiple E2 motifs within E6 promoter (p89). Thus, an assay was performed to determine whether AAV Rep78 might bind to a sequence (nt 7758-7930) from p89 which contains two such motifs in close proximity. The EMSA results, shown in Figure 9A demonstrate that AAV Rep78 is able to recognize and bind E2 motif DNA from BPV-1 p89. Figure 9B shows that, as with the AAV Rep78-p97 interaction, both the unlabeled BPV-1 p89 and the AAV TR DNAs were able to compete against AAV Rep78-³²P-BPV-1 LCR interaction, however TR DNA was the better competitor.

AAV Rep78 mutants effect on the AAV p5 promoter

Figure 10 provides an example of AAV Rep78 mutants, Rep-77^{LG} and Rep-79^{FA} that are unable to bind to the AAV p5 promoter and unable to inhibit the p5 promoter by *in*

vitro transcription. The promoter, p5, is an important promoter as it expresses AAV Rep78 itself, which is critical for AAV replication. This mutant also is not able to bind to HPV-16 p97. The present invention shows that this inability to bind results in higher levels of AAV genome replication and virion production. The Rep-192^{HG} is able to inhibit p5 (AAV promoter) better than wild-type AAV Rep78, Rep-77^{LG} and Rep79^{FA}, which do not inhibit p5 at all. All data indicates that the mutant inhibition of AAV p5, HPV-16 p97 and HIV-1 long terminal repeat are similar. Additionally, the level of p5 transcription mimics and supports the level of DNA replication of these mutants as FLAGS for wild-type 620, Rep-77^{LG} and Rep-192^{HG}.

The present inventor discovered that by including this particular mutation in a full length AAV genome (FLAG-77^{LG}), he was able to obtain replication levels of 150% above wild-type AAV-2, which makes it useful as a complementor for generating recombinant AAV. Figure 11 shows that FLAG-77^{LG} replicates at higher levels and FLAG-192^{HG} replicates at slightly depressed levels compared to wild-type AAV Rep78 (pSM620/Sph). Although FLAG-192^{HG} does not show a very low replication level in the experiment shown in Figure 11 but over 5 experiments, FLAG-192^{HG} clearly replicates at lower levels than wild type pSM620. Another AAV Rep78 mutant that is unable to bind to p97 is Rep-64^{LH}65TM.

An example of an AAV Rep78 mutant that has enhanced binding as well as different binding to the AAV TR (terminal repeat) and the p5 is the Rep-192^{HG} as shown by a higher sized band on the EMSA in Figure 12 and Figure 13 as compared to the wild-type AAV Rep78 (MBP-Rep78), which makes it useful for anti-cancer protocols of treatment. Figure 12 shows that Rep192^{HG} binds TR DNA in a different higher order complex and with higher affinity. The FLAG-192^{HG} replicates at lower levels than wild-type AAV Rep78, which is consistent with increased self-inhibition. AAV Rep192^{HG} is a super binder, and therefore, a super inhibitor, as shown by the higher and darker band in the Rep192^{HG} lane representing a protein-DNA complex.

The present invention discloses that the AAV Rep78 protein meaningfully binds to the p97 promoter of HPV-16. This binding, although not as strong as Rep78-AAV TR interaction, is clearly above the so-called "non-specific" DNA-binding ability of Rep78 as determined by the experiments presented in Figure 1. The importance of this interaction is supported by the finding that two AAV mutants (FLAG-64^{LH}65TM and FLAG-77^{LG}), whose

Rep78 proteins are unable to recognize the HPV-16 LCR p97 target, are defective for *in vitro* transcriptional inhibition of p97 and for the inhibition of HPV-16 directed oncogenic transformation. It should also be noted that Rep78 is unable to significantly affect expression from the MSV-LTR (18) and Rep78 is also unable to bind partial (Batchu, *et al.*, *Cancer Letters* 86:23-31 (1994), Kokorina, *et al.*, *J. Hum. Virol* 1:441-450 (1998)) or
5 the full length sequence of this promoter (Figure 1B).

Figure 14 shows the inhibition of HPV-16 by AAV Rep78 mutants. pL67R (HPV16+ras) and each of the indicated AAV plasmids were cotransfected (4 ug each) into contact inhibited C127 cells and allowed to grow for 2 1/2 weeks, and oncogenic foci
10 counted. Note that inhibition of HPV-16-induced oncogenic foci roughly matches the DNA binding ability of the Rep78 protein. The results show that Rep-192^{IG} inhibits HPV-16 oncogenic transformation and p97 promoter expression better than 620sph (wild-type) Rep 78, which demonstrates the anti-oncogenic properties of this Rep78 mutant.

One previous relevant study undertaken by Horer *et al.*, *J. Virol.* 69:5485-5496
15 (1995) attempted to identify the *cis*-responsible element within the analogous p105 promoter of HPV-18. Their results, obtained by the deletion and mutation of sequences along the length of the p105 promoter, were inconclusive for finding a specific responsible element. Their interpretation of this data was that the mechanism of inhibition was complex and involved multiple components. The data disclosed herein may suggest such
20 complexity, for although the phenotypes of mutants Rep-64^{LH}65TM and Rep-77^{LG} are quite strong in Figures 5, it is also clear that they are still able to mildly inhibit oncogenic transformation in Figure 7 (only small foci were generated). It is reasonable to suggest that Rep78's protein-protein interactions are likely also involved in these other inhibitory pathways. In addition to Rep78's ability to meaningfully interact with transcription factors,
25 in preliminary experiments, Rep78-E7 oncoprotein interaction by both Western blot, affinity chromatography, and yeast GAL4 two-hybrid cDNA analyses has been observed (Zhan and Hermonat, unpublished).

The fact that Rep78 binds HPV-16 p97 is surprising as this target sequence contains no GAGC motifs, the core sequence of most Rep78 DNA recognition. Rep78 binding to
30 promoter sequences has been observed before, including the AAV p5 promoter (McCarty, *et al.*, *J. Virol.* 74:4988-4997 (1994)), the c-H-ras promoter (Batchu, *et al.*, *Cancer Letters* 86:23-31 (1994)), the human immunodeficiency virus long terminal repeat TAR region

(Batchu, *et al.*, *FEBS Letters* 367:267-271 (1995), Kokorina, *et al.*, *J. Hum. Virol* 1:441-450 (1998)), and the cytomegalovirus immediate early promoter (Wonderling, *et al.*, *J. Virol.* 71:2528-2534 (1996)). All of these promoters have GAGC (or GCTC) motifs, and binding by the large Rep proteins (proteins 78/68) appears to be required for the inhibition of the AAV p5 promoter (Kyostio, *et al.*, *J. Virol.* 69:6787-6796 (1995)) and the HIV-LTR (Kokorina, *et al.*, *J. Hum. Virol* 1:441-450 (1998)). In experiments using Rep78 affinity selection of random DNA sequences the consensus target sequence contained a duplex GAGC motif (Chiorini, *et al.*, *J. Virol.* 69:7334-7338 (1995)). However a small subset of these selected sequences had no GAGC motifs. Thus, there is a precedence for Rep78 binding DNA without GAGC motifs. However, none of these sequences have significant homology with p97. Furthermore, none of the random selected sequences contain interrupted palindromes as are present in p97. It has also been observed that Rep78 binds a region within the BPV-1 LCR, which contains two E2 motifs, and it is believed that these motifs are the specific target for Rep78 recognition. AAV's own TRs, Rep78's natural and favored substrate, also have significant secondary structure. This further strengthens the argument that such structures (E2 or otherwise) are involved in Rep78 recognition of p97. However, the phenotype of Rep-77^{LG} suggests that Rep78's recognition of AAV TR DNA is different than its recognition of p97. The finding that Rep78 specifically recognizes the negative strand of the p97 promoter is novel (Figure 3). These data further indicate that Rep78 discriminates between single stranded DNA substrates, by sequence, as it does for double stranded substrates. Finally, Rep78 binding to p97 may also affect the E1-E2 complex from binding the origin of replication which is located just upstream (Sedman, *et al.* *EMBO J.* 15:5085-5092, 1996 (1996), Yasugi, *et al.*, *J. Virol.* 71:891-899 (1997), Sarafi, *et al.*, *Virology* 211:385-396 (1995)). Rep78 is known to inhibit BPV-1 DNA replication (Hermonat, P.L. *Virology* 189:329-333 (1992)). Rep 79^{FA} has a similar phenotype to Rep77^{LG} in that it does not bind to the p5 and p97 promoter sequences very well but unlike Rep77^{LG}, Rep79^{FA} does not bind to TR DNA.

All of the publications cited herein are incorporated in their entirety by reference as is an article entitled "Binding of the human papillomavirus type 16 p97 promoter by the adeno-associated virus Rep78 major regulatory protein correlates with inhibition" by Zhan *et al.*, *J. Biol. Chem.* 274: 31619-31624 (1999).

WE CLAIM:

1. An AAV Rep78 mutant comprising an AAV Rep78 modified protein that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein.
2. The AAV Rep78 mutant of claim 1, wherein said AAV Rep78 modified protein binds to at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, an oncogene or a HIV differently as compared to the binding of said wild-type AAV Rep78 protein.
3. The AAV Rep78 mutant of claim 2, wherein said different DNA binding is selected from the group consisting of no DNA binding, weak DNA binding and enhanced DNA binding as compared to the binding of said wild-type AAV Rep78 protein.
4. The AAV Rep78 mutant of claim 3, wherein said mutant having no DNA binding or weak DNA binding to said DNA sequence obtained from at least one of a papillomavirus, an AAV, an oncogene or a HIV that results in the generation of higher levels of AAV DNA replication and virion numbers.
5. The AAV Rep78 mutant of claim 3, wherein said mutant having enhanced DNA binding to said DNA sequence obtained from at least one of a papillomavirus or an oncogene that results in enhanced inhibition of at least one of a papillomavirus or an oncoprotein.
6. The AAV Rep78 mutant of any one of claims 2-5, wherein said mutant is selected from the group consisting of a truncated wild-type AAV Rep78 protein, a wild-type AAV Rep78 protein containing amino acid substitutions, a wild-type AAV Rep78 protein containing internal amino acid deletions, and a combination thereof.
7. The AAV Rep78 mutant of claim 6, wherein said mutant is a truncated AAV Rep78 protein containing at least the minimum number of amino acids of the wild-type AAV Rep78 protein necessary to bind to said DNA sequence to obtain enhanced inhibition of a papillomavirus or an oncogene.

8. The AAV Rep78 mutant of any one of claims 2-7, wherein said DNA sequence to which said mutant binds is a promoter region of said papillomavirus, said AAV or said oncogene.
9. The AAV Rep78 mutant of claim 8, wherein said papillomavirus promoter region is nucleotides 14-56 of p97 of HPV-16.
10. The AAV Rep78 mutant of claim 7, comprising at least two truncated wild-type AAV Rep78 linked to form a multimer AAV Rep78 mutant.
11. The AAV Rep78 mutant of claim 4, wherein said mutant is AAV Rep-77^{LG}, AAV Rep-79^{FA}.
12. The AAV Rep78 mutant of claim 5, wherein said mutant is AAV Rep-192^{HG}.
13. A fusion protein comprising the AAV Rep78 protein or said AAV Rep78 mutant of any one of claims 1-12.
14. The fusion protein of claim 13, further comprising the *tat* protein of HIV linked to said AAV Rep78 mutant.
15. The fusion protein of claim 14, wherein said *tat* protein is the *tat* protein of HIV-1.
16. The fusion protein of any one of claims 13-15, further comprising a maltose-binding protein (MBP) linked to said AAV Rep78 protein or said AAV Rep78 mutant.
17. A pharmaceutical composition comprising at least one AAV Rep78 mutant or said AAV Rep78 protein according to any one of claims 1-16, in admixture with a pharmaceutically acceptable carrier.
18. A method of treating papillomavirus-associated diseases, cancer or HIV-associated diseases comprising administering said pharmaceutical composition of claim 17 to a patient afflicted with a papillomavirus associated disease or cancer.

19. A DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-16.
20. The DNA construct comprising said DNA sequence of claim 19 and a vector.
21. The DNA construct of claim 20 wherein said vector is selected from the group consisting of an AAV, a rAAV, a retrovirus, an adenovirus and a liposome.
22. The DNA construct of any one of claims 19-21, further comprising an inducible promoter operably linked to said DNA sequence.
23. The DNA construct of claim 22, wherein said inducible promoter is controlled by a viral protein or an oncoprotein.
24. A pharmaceutical composition comprising at least one of the DNA sequences of any one of claims 19-23 in admixture with a pharmaceutically acceptable carrier.
25. A method of inhibiting papillomavirus associated diseases or cancer comprising administering said pharmaceutical composition of claim 24 to a patient afflicted with a papillomavirus associated disease or cancer.
26. The method of claim 25, wherein said inducible promoter is controlled by the expression of a papillomavirus protein or an oncoprotein in said patient as a result of said disease or cancer.
27. A DNA sequence comprising the full length AAV genome modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 19 to 23.
28. A DNA sequence comprising at least the minimum portion of the AAV genome sufficient to complement a defective rAAV, wherein said portion of the AAV genome is modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 19 to 23.

29. A method of producing recombinant AAV (rAAV) comprising transducing a susceptible mammalian cell with a DNA sequence of claims 27 or 28 and a DNA sequence encoding said rAAV to obtain rAAV virions.

30. The method of claim 29, wherein said DNA sequence is sufficient to complement a defective rAAV comprises at least the AAV *lip-cap* gene and said DNA sequence encoding an AAV Rep78 mutant.

31. The method of claim 30, wherein said rAAV comprises a DNA sequence encoding a heterologous gene.

32. A method of treating papillomavirus associated diseases or cancer comprising administering a pharmaceutical composition comprising a AAV Rep78 protein to a patient afflicted with a papillomavirus associated disease or cancer, wherein said AAV Rep78 protein binds to a papillomavirus promoter that controls the expression of a papillomavirus oncoprotein or binds to a promoter that controls the expression of an oncoprotein.

33. The method of claim 32, wherein said AAV Rep78 binds to nucleotides 14-56 of p97 of HPV-16 inhibiting expression of HPV.

34. An AAV Rep78 regulation element comprising the nucleotides shown in the nucleotide sequence of Figure 2, wherein said element provides the binding site for the AAV Rep78 protein.

35. The AAV Rep78 regulation element of claim 34 comprising about nucleotides 14 -56 of the nucleotide sequence of Figure 2.

36. An AAV Rep78 regulatable promoter comprising the regulation element of claims 34 or 35 and the remaining promoter sequences from a promoter other than the HPV-16 p97 promoter.


37. A method of selecting an AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein, wherein said method comprises:


contacting an AAV Rep78 mutant of any one of claims 1-16 with at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, a HIV or an oncogene for a period of time to allow binding of said mutant to said DNA; and

determining said binding of said mutant to select said mutant that binds differently to said DNA than said wild-type Rep78 protein.

38. Use of the pharmaceutical composition of claims 17 or 24 for the treatment of a papillomavirus-associated disease, cancer or a HIV-associated diseases.

39. Use of the AAV Rep78 mutant or the AAV Rep78 protein of any one of claims 1-16 or the DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-16 to prepare a pharmaceutical composition for the treatment of a papillomavirus-associated diseases, cancer or HIV-associated diseases



**FIG. 2**

3/18






FIG. 3

4/18

**FIG. 4**

5/18

**FIG. 5**

6/18




FIG. 6

7/18




FIG. 7

A		B			
cold compet	TR	p97	p97	TR	p97
DNA (ug)	0	0	0	cold compet	
MBP-Rep78	-	+	+	DNA (ug) 0	
Rep-77 ^{LG}	-	-	-	MBP-Rep78	+
³² P-TR	+	+	+	³² P-p97	+




FIG. 8

9/18




FIG. 9

10/18






FIG. 10

11/18

**FIG. 11**

12/18

**FIG. 12**

13/18




FIG. 13

14/18

plasmids transfected	oncogenic foci
pL67R + 334 (no Rep)	34
pL67R + 620/Sph (wild type)	8
pL67R + Rep77 ^{LG}	19
pL67R + Rep79 ^{FA}	25
pL67R.+ Rep192 ^{HG}	4

FIG. 14

15/18

FIG. 15A

```

1  ttggccactc cctctctgcg cgctcgctcg ctactgagg ccgggcgacc aaaggctcgcc
61  cgacgcccgg gctttgcccc ggcggcctca gtgagcgagc gagcgcgag agagggagtg
121  gccaaactca tcactagggg ttcttgagg ggtggagtcg tgacgtgaat tacgtcatag
181  ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacacccat
241  gtggtcacgc tgggtattta agcccagtg agcacgcagg gtctccattt tgaagcggga
301  ggtttgaacg cgcagccgcc atgcccgggt ttacgagat tggattaaag gtccccagcg
361  accttgacgg gcatctgccc ggcatttctg acaqctttgt gaaactgggtg gccgagaaag
421  aatgggagtt gccgccagat tctgacatgg atctgaatct gattgagcag gcaccctga
481  ccgtggccga gaagctgcag cgcgactttc tgacggaaatg gcgccgtgtg agtaaggccc
541  cgagggccct tttctttgtg caatttgaga agggagagag ctacttccac atgcacgtgc
601  tcgtggaaac caccggggtg aaatccatgg ttttgggacg tttcctgagt cagattcgcg
661  aaaaactgat tcagagaatt taccgcggga tcgagccgac tttgccaaac tggttcgcg
721  tcacaaagac cagaaatggc gccggaggcg ggaacaaagt ggtggatgag tgctacatcc
781  ccaattactt gctccccaaa acccagcctg agctccagt ggcgtggact aatatggaac
841  agtatttaag cgcctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga
901  cgcacgtgtc gcagacgcag gacgagaaca aagagaatca gaatcccaat tctgatgcgc
961  cgtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca
1021 aggggattac ctcgagaaag cagtggatcc aggaggacca ggcctcatat atctccttca
1081 atgcggcctc caactcgcgg tcccaaatca aggcctgcctt ggacaaatgcg ggaagatta
1141 tgagcctgac taaaaccgcc cccgactacc tgggtgggcca gcagcccgtg gaggacattt
1201 ccagcaatcg gatttataaa attttggaac taaacgggta cgatccccc aaatgcggctt
1261 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg
1321 ggcctgcaac taccgggaag accaacatcg cggaggccat agcccacact gtgcccttct
1381 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg
1441 tgatctggtg ggaggagggg agatgaccg ccaaggtcgt ggagtcggcc aaagccattc
1501 tcggagggaag caaggtgcgc gtggaccaga aatgcaagtc ctcggcccag atagaccga
1561 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga
1621 ccttcgaaca ccagcagccg ttgcaagacc ggatgttcaa atttgaactc accgcgcgtc

```

SUBSTITUTE SHEET (RULE 26)

16/18

1681 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cgtgggcaa
 1741 aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaagggtgga gccaagaaaa
 1801 gaccgcccc cagtgcgca gataaagtg agccaaacg ggtgcgcgag tcagttgcgc
 1861 agccatcgac gtcagacgc gaagcttgcg tcaactacgc agacaggtac caaaacaaat
 1921 gttctcgtca cgtgggcatg aatctgatgc tgttccctg cagacaaatgc gagagaatga
 1981 atcagaattc aaatatctgc ttactcacg gacagaaaga ctgttttagag tgctttccc
 2041 tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgtg tcagaaactg tgctacattc
 2101 atcatatcat gggaaggtg ccagacgctt gcactgcctg cgatctggtc aatgtggatt
 2161 tggatgactg catctttgaa caataaatga tttaaatcag gtatggctgc cgaatggtat
 2221 cttccagatt ggctcgagga cactctctct gaaggaataa gacagtgtgtg gaagctcaaa
 2281 cctggccac caccaccaa gcccgagag cggcataaag acgacagcag ggtcttgtg
 2341 cttcctgggt acaagtacct cggacccttc aacggactcg acaaggaga gccggtcaac
 2401 gaggcagacg ccgcccct tacctcaagt acaaccacgc caagcctac gaccggcagc tcgacagcgg
 2461 agacaacccg tacctcaagt acaaccacgc cgacgcggag tttagaggc gccttaaga
 2521 agatacgtct tttgggggca acctcggaag agcagctctc caggcgaaa agagggtctt
 2581 tgaacctctg ggcctgggtg agaacctgt taagacggct ccgggaaaa agaggccggt
 2641 agagcactct cctgtggagc cagactcctc ctgggaacc ggaaggcgg gccagcagcc
 2701 tgaagaaaa agattgaatt ttggtcagac tgagacgca gactcagtac ctgaccccca
 2761 gcctctcgga cagccaccag cagccccctc tgggtcggg actaatcga tggctacagg
 2821 cagtggcgca ccaatggcag acaataacga gggcgcgcgac ggagtgggtg attcctccgg
 2881 aaattggcat tgcgattcca catggatggg cgacagagtc atcaccacca gacccgaaac
 2941 ctgggcccctg ccacacctaca acaaccacct ctacaaaca atttccagcc aatcaggagc
 3001 ctcgaacgac aatcactact ttggctacag cacccttgg ggtattttg acttcaacag
 3061 attccactgc cacttttcac cactgactg gcaagactc atcaacaaca actggggatt
 3121 ccgaaccaag agactcaact tcaagctctt taacattcaa gtcaagagg tcacgcagaa
 3181 tgacgggtacg acgacgattg ccaataacct taccagcagc gttcaggtgt ttactgactc
 3241 ggagtaccag ctcccgtacg tcctcggctc ggcgcataca ggatgcctcc cgcggttccc
 3301 agcagacgct ttcattggtc cacagtatgg atacctcacc ctgaacaacg ggagtcaggc

FIG. 15B

17/18

3361 agtaggacgc tcttcatttt actgcctgga gtactttcct tctcagatgc tgcgtaccgg
 3421 aaacaacttt accttcagct acacttttga ggacgttcct ttcacagca gctacgctca
 3481 cagccagagt ctggaccgtc tcatgaatcc tctcatcgac cagtacctgt attacttgag
 3541 cagaacaaac actccaagtg gaaccaccac gcagtcaagg cttcagtttt ctcaggcccgg
 3601 agcgagtgc attcgggacc agtctaggaa ctggcttcct ggacctgtt accgccagca
 3661 gcgagtatca aagacatctg cggataacaa caacagtga tactcgtgga ctggagctac
 3721 caagtaccac ctcaatggca gagactctct ggtgaatccg gccatggcaa gccacaagga
 3781 cgatgaagaa aagttttttc ctcagagcgg ggttctcatc ttggggaagc aaggctcaga
 3841 gaaaacaaat gtgaacattg aaaaggctcat gattacagac gaagaggaaa tcggaacaac
 3901 caatcccctq gctacggagc agtatggttc tgtatctacc aacctccaga gaggcaacag
 3961 acaagcagct accgcagatg tcaacacaca aggcgttctt ccaggcatgg tctggcagga
 4021 cagagatgtg taccttcagg ggcccatctg ggcaaaagatt ccacacacgg acggacattt
 4081 tcacccctct cccctcatgg gtggattcgg acttaaacac cctcctccac agattctcat
 4141 caagaacacc cgggtacctg cgaatccttc gaccaccttc agtgcggcaa agtttgcttc
 4201 ctctcatcaca cagtactcca cgggacacgg tcagcgtgga gatcagatgg gagctgcaga
 4261 aggaaaaacag caaacgctgg aatcccga aaatccgtacac ttccaactac aacaagtctg
 4321 ttaatcgtgg acttacctg gatactaag gcgtgtattc agagcctcgc cccattggca
 4381 ccagataacct gactcgtaat ctgtaattgc ttgttaatca ataaaccgtt taattcgttt
 4441 cagttgaact ttggtctctg cgtatttctt tcttatctag ttccatggc tacgtagata
 4501 agtagcatgg cgggttaatc attaaactaca aggaaccctt agtgatggag ttggccactc
 4561 cctctctgcg cgctcgctcg ctcaactgag ccgggcgacc aaaggtcgcc cgacgcccgg
 4621 gctttgcccg ggcggcctca gtgagcgagc ggcgcgcag agaggagtg gccaa

FIG. 15C

18/18

MPGFYEIVIKVPSDL DGHLPGISDSFVNWVAEKEWELPPDSMD
LNLIEQAPLTVAEKLQRDFLTEWRRVSKAPEALFFVQFEKGESYFHMHVLVETTGVS
MVLGRFLSQIREKLIQRIYRGIEPTLPNWFAVTKTRNGAGGNKVVD ECIYPNYLLPK
TQPELQWAWTNMEQYLSACLNLTERKRLVAQHLTHVSQTQEKNENQNPNSDAPVIRS
KTSARYMELVGWLVDKGITSEKQWIQEDQASYISFNAA SNRSQIKAALDNAGKIMSL
TKTAPDYLVGQPVEDISSNRIYKILELNGYDPQYAA SVFLGWATKKFGKRNTIWLFG
PATTGKTNIAEIAHTVPFYGCCVNW TNENFPFND CVDKMVIWEEGKMTAKVVESAKA
ILGGSKVRVDQCKSSAQIDPTPVI VTSNTNMCAVIDGNSTTFEHQQPLQDRMFKFEL
TRRLDHDGKVTKQEVKDFFRWAKDHVVEVEHEFYVKKGGAKKR PAPSADADISEPKRV
RESVAQPSTSDAEASINYADRYQNKCSRHVGMNLM L FPCRQ CERMNQNSNICFTHGQK
DCLECFPVSESQPVSVVKKAYQKLCYIHHIMGKV PDACTACDLVNVDLDDCIFEQ

FIG. 16


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 May 2001 (10.05.2001)

PCT

(10) International Publication Number
WO 01/32711 A3

- (51) International Patent Classification⁷: **A61K 48/00**,
C07K 14/075, C12N 15/86
- (21) International Application Number: **PCT/US00/29187**
- (22) International Filing Date: 23 October 2000 (23.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/160,608 21 October 1999 (21.10.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/160,608 (CIP)
Filed on 21 October 1999 (21.10.1999)
- (71) Applicant (for all designated States except US):
BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS [US/US]; 2404 North University Avenue, Little Rock, AR 72207-3608 (US).
- (72) Inventor; and
(75) Inventor/Applicant (for US only): **HERMONAT, Paul**, L. [US/US]; 1812 Reservoir Road, Apartment 143, Little Rock, AR 72227 (US).
- (74) Agent: **GRANADOS, Patricia, D.**; Foley & Lardner, 3000 K. Street, N.W., Suite 500, Washington, DC 20007-5109 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **ADENO-ASSOCIATED VIRUS AAV REP78 MAJOR REGULATORY PROTEIN, MUTANTS THEREOF AND USES THEREOF**

(57) Abstract: AAV Rep78 mutants comprising a modified AAV Rep78 protein that possess different biochemical and biological functions as compared to the wild-type AAV Rep78 protein are disclosed. Particularly, the AAV Rep78 mutants that bind to at least one of a papillomavirus DNA or an AAV DNA or an oncogene or HIV DNA differently as compared to the wild-type AAV Rep78 protein. The present invention additionally is directed to a method of inhibiting papillomavirus associated diseases comprising administering pharmaceutical compositions containing AAV Rep78 mutants or the DNA sequences encoding the mutants. Also disclosed are DNA sequences comprising the full length AAV genome modified to where the AAV Rep78 protein is replaced with an AAV Rep mutant that binds weakly or not at all to a papillomavirus DNA or an AAV DNA or both differently as compared to the wild-type AAV Rep78 protein.

WO 01/32711 A3



Published:

- with international search report
- with amended claims

Date of publication of the amended claims: 24 January 2002

(88) Date of publication of the international search report:
27 December 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

AMENDED CLAIMS

[received by the International Bureau on 21 September 2001 (21.09.01);
original claims 1-39 replaced by amended claims 1-37 (5 pages)]

1. An AAV Rep78 mutant comprising an AAV Rep78 modified protein that binds to at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, an oncogene or a HIV differently as compared to the binding of said wild-type AAV Rep78 protein, and that results in AAV DNA replication and/or AAV virion production, wherein said different DNA binding is selected from the group consisting of no DNA binding, weak DNA binding and enhanced DNA binding as compared to the binding of said wild-type AAV Rep78 protein.
2. The AAV Rep78 mutant of claim 1, wherein said mutant having no DNA binding or weak DNA binding to said DNA sequence obtained from at least one of a papillomavirus, an AAV, an oncogene or a HIV that results in the generation of higher levels of AAV DNA replication and virion numbers as compared to the wild-type Rep78.
3. The AAV Rep78 mutant of claim 1, wherein said mutant having enhanced DNA binding to said DNA sequence obtained from at least one of a papillomavirus or an oncogene that results in enhanced inhibition of at least one of a papillomavirus or an oncoprotein.
4. The AAV Rep78 mutant of any one of claims 1-3, wherein said mutant is selected from the group consisting of a truncated wild-type AAV Rep78 protein, a wild-type AAV Rep78 protein containing amino acid substitutions, a wild-type AAV Rep78 protein containing internal amino acid deletions, and a combination thereof.
5. The AAV Rep78 mutant of claim 4, wherein said mutant is a truncated AAV Rep78 protein containing at least the minimum number of amino acids of the wild-type AAV Rep78 protein necessary to bind to said DNA sequence to obtain enhanced inhibition of a papillomavirus or an oncogene.

6. The AAV Rep78 mutant of any one of claims 1-5, wherein said DNA sequence to which said mutant binds is a promoter region of said papillomavirus, said AAV or said oncogene.
7. The AAV Rep78 mutant of claim 6, wherein said papillomavirus promoter region is nucleotides 14-56 of p97 of HPV-16.
8. The AAV Rep78 mutant of claim 5, comprising at least two truncated wild-type AAV Rep78 linked to form a multimer AAV Rep78 mutant.
9. The AAV Rep78 mutant of claim 2, wherein said mutant is AAV Rep-77^{LG}, AAV Rep-79^{FA}.
10. The AAV Rep78 mutant of claim 3, wherein said mutant is AAV Rep-192^{HG}.
11. A fusion protein comprising the AAV Rep78 protein or said AAV Rep78 mutant of any one of claims 1-10.
12. The fusion protein of claim 11, further comprising the *tat* protein of HIV linked to said AAV Rep78 mutant.
13. The fusion protein of claim 12, wherein said *tat* protein is the *tat* protein of HIV-1.
14. The fusion protein of any one of claims 11-13, further comprising a maltose-binding protein (MBP) linked to said AAV Rep78 protein or said AAV Rep78 mutant.
15. A pharmaceutical composition comprising at least one AAV Rep78 mutant or said AAV Rep78 protein according to any one of claims 1-14, in admixture with a pharmaceutically acceptable carrier.

SUBSTITUTE PAGE

16. A method of treating papillomavirus-associated diseases, cancer or HIV-associated diseases comprising administering said pharmaceutical composition of claim 15 to a patient afflicted with a papillomavirus associated disease or cancer.
17. A DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-14.
18. The DNA construct comprising said DNA sequence of claim 17 and a vector.
19. The DNA construct of claim 18, wherein said vector is selected from the group consisting of an AAV, a rAAV, a retrovirus, an adenovirus and a liposome.
20. The DNA construct of any one of claims 17-19, further comprising an inducible promoter operably linked to said DNA sequence.
21. The DNA construct of claim 20, wherein said inducible promoter is controlled by a viral protein or an oncoprotein.
22. A pharmaceutical composition comprising at least one of the DNA sequences of any one of claims 17-21 in admixture with a pharmaceutically acceptable carrier.
23. A method of inhibiting papillomavirus associated diseases or cancer comprising administering said pharmaceutical composition of claim 22 to a patient afflicted with a papillomavirus associated disease or cancer.
24. The method of claim 22, wherein said inducible promoter is controlled by the expression of a papillomavirus protein or an oncoprotein in said patient as a result of said disease or cancer.

25. A DNA sequence comprising the full length AAV genome modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 17 to 21.
26. A DNA sequence comprising at least the minimum portion of the AAV genome sufficient to complement a defective rAAV, wherein said portion of the AAV genome is modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 17 to 21.
27. A method of producing recombinant AAV (rAAV) comprising transducing a susceptible mammalian cell with a DNA sequence of claims 25 or 26 and a DNA sequence encoding said rAAV to obtain rAAV virions.
28. The method of claim 27, wherein said DNA sequence is sufficient to complement a defective rAAV comprises at least the AAV *lip-cap* gene and said DNA sequence encoding an AAV Rep78 mutant.
29. The method of claim 28, wherein said rAAV comprises a DNA sequence encoding a heterologous gene.
30. A method of treating papillomavirus associated diseases or cancer comprising administering a pharmaceutical composition comprising a AAV Rep78 protein to a patient afflicted with a papillomavirus associated disease or cancer, wherein said AAV Rep78 protein binds to a papillomavirus promoter that controls the expression of a papillomavirus oncoprotein or binds to a promoter that controls the expression of an oncoprotein.
31. The method of claim 30, wherein said AAV Rep78 binds to nucleotides 14-56 of p97 of HPV-16 inhibiting expression of HPV.

32. An AAV Rep78 regulation element comprising the nucleotides shown in the nucleotide sequence of Figure 2, wherein said element provides the binding site for the AAV Rep78 protein.

33. The AAV Rep78 regulation element of claim 32 comprising about nucleotides 14 -56 of the nucleotide sequence of Figure 2.

34. An AAV Rep78 regulatable promoter comprising the regulation element of claims 32 or 33 and the remaining promoter sequences from a promoter other than the HPV-16 p97 promoter.

35. A method of selecting an AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein, wherein said method comprises:

contacting an AAV Rep78 mutant of any one of claims 1-14 with at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, a HIV or an oncogene for a period of time to allow binding of said mutant to said DNA; and

determining said binding of said mutant to select said mutant that binds differently to said DNA than said wild-type Rep78 protein.

36. Use of the pharmaceutical composition of claims 15 or 22 for the treatment of a papillomavirus-associated disease, cancer or a HIV-associated diseases.

37. Use of the AAV Rep78 mutant or the AAV Rep78 protein of any one of claims 1-14 or the DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-14 to prepare a pharmaceutical composition for the treatment of a papillomavirus-associated diseases, cancer or HIV-associated diseases

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2001 (10.05.2001)


PCT

(10) International Publication Number
WO 01/32711 A3

- (51) International Patent Classification⁷: **A61K 48/00**,
C07K 14/075, C12N 15/86
- (21) International Application Number: **PCT/US00/29187**
- (22) International Filing Date: 23 October 2000 (23.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/160,608 21 October 1999 (21.10.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/160,608 (CIP)
Filed on 21 October 1999 (21.10.1999)
- (71) Applicant (for all designated States except US):
BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS [US/US]; 2404 North University Avenue, Little Rock, AR 72207-3608 (US).
- (72) Inventor; and
(75) Inventor/Applicant (for US only): **HERMONAT, Paul, L.** [US/US]; 1812 Reservoir Road, Apartment 143, Little Rock, AR 72227 (US).
- (74) Agent: **GRANADOS, Patricia, D.;** Foley & Lardner, 3000 K. Street, N.W., Suite 500, Washington, DC 20007-5109 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: **ADENO-ASSOCIATED VIRUS AAV REP78 MAJOR REGULATORY PROTEIN, MUTANTS THEREOF AND USES THEREOF**



(57) Abstract: AAV Rep78 mutants comprising a modified AAV Rep78 protein that possess different biochemical and biological functions as compared to the wild-type AAV Rep78 protein are disclosed. Particularly, the AAV Rep78 mutants that bind to at least one of a papillomavirus DNA or an AAV DNA or an oncogene or HIV DNA differently as compared to the wild-type AAV Rep78 protein. The present invention additionally is directed to a method of inhibiting papillomavirus associated diseases comprising administering pharmaceutical compositions containing AAV Rep78 mutants or the DNA sequences encoding the mutants. Also disclosed are DNA sequences comprising the full length AAV genome modified to where the AAV Rep78 protein is replaced with an AAV Rep mutant that binds weakly or not at all to a papillomavirus DNA or an AAV DNA or both differently as compared to the wild-type AAV Rep78 protein.

WO 01/32711 A3



Published:

- with international search report
- with amended claims

Date of publication of the amended claims: 24 January 2002

(88) Date of publication of the international search report:
27 December 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

AMENDED CLAIMS

[received by the International Bureau on 21 September 2001 (21.09.01);
original claims 1-39 replaced by amended claims 1-37 (5 pages)]

1. An AAV Rep78 mutant comprising an AAV Rep78 modified protein that binds to at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, an oncogene or a HIV differently as compared to the binding of said wild-type AAV Rep78 protein, and that results in AAV DNA replication and/or AAV virion production, wherein said different DNA binding is selected from the group consisting of no DNA binding, weak DNA binding and enhanced DNA binding as compared to the binding of said wild-type AAV Rep78 protein.
2. The AAV Rep78 mutant of claim 1, wherein said mutant having no DNA binding or weak DNA binding to said DNA sequence obtained from at least one of a papillomavirus, an AAV, an oncogene or a HIV that results in the generation of higher levels of AAV DNA replication and virion numbers as compared to the wild-type Rep78.
3. The AAV Rep78 mutant of claim 1, wherein said mutant having enhanced DNA binding to said DNA sequence obtained from at least one of a papillomavirus or an oncogene that results in enhanced inhibition of at least one of a papillomavirus or an oncoprotein.
4. The AAV Rep78 mutant of any one of claims 1-3, wherein said mutant is selected from the group consisting of a truncated wild-type AAV Rep78 protein, a wild-type AAV Rep78 protein containing amino acid substitutions, a wild-type AAV Rep78 protein containing internal amino acid deletions, and a combination thereof.
5. The AAV Rep78 mutant of claim 4, wherein said mutant is a truncated AAV Rep78 protein containing at least the minimum number of amino acids of the wild-type AAV Rep78 protein necessary to bind to said DNA sequence to obtain enhanced inhibition of a papillomavirus or an oncogene.

6. The AAV Rep78 mutant of any one of claims 1-5, wherein said DNA sequence to which said mutant binds is a promoter region of said papillomavirus, said AAV or said oncogene.
7. The AAV Rep78 mutant of claim 6, wherein said papillomavirus promoter region is nucleotides 14-56 of p97 of HPV-16.
8. The AAV Rep78 mutant of claim 5, comprising at least two truncated wild-type AAV Rep78 linked to form a multimer AAV Rep78 mutant.
9. The AAV Rep78 mutant of claim 2, wherein said mutant is AAV Rep-77^{LG}, AAV Rep-79^{FA}.
10. The AAV Rep78 mutant of claim 3, wherein said mutant is AAV Rep-192^{HG}.
11. A fusion protein comprising the AAV Rep78 protein or said AAV Rep78 mutant of any one of claims 1-10.
12. The fusion protein of claim 11, further comprising the *tat* protein of HIV linked to said AAV Rep78 mutant.
13. The fusion protein of claim 12, wherein said *tat* protein is the *tat* protein of HIV-1.
14. The fusion protein of any one of claims 11-13, further comprising a maltose-binding protein (MBP) linked to said AAV Rep78 protein or said AAV Rep78 mutant.
15. A pharmaceutical composition comprising at least one AAV Rep78 mutant or said AAV Rep78 protein according to any one of claims 1-14, in admixture with a pharmaceutically acceptable carrier.

SUBSTITUTE PAGE

16. A method of treating papillomavirus-associated diseases, cancer or HIV-associated diseases comprising administering said pharmaceutical composition of claim 15 to a patient afflicted with a papillomavirus associated disease or cancer.
17. A DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-14.
18. The DNA construct comprising said DNA sequence of claim 17 and a vector.
19. The DNA construct of claim 18, wherein said vector is selected from the group consisting of an AAV, a rAAV, a retrovirus, an adenovirus and a liposome.
20. The DNA construct of any one of claims 17-19, further comprising an inducible promoter operably linked to said DNA sequence.
21. The DNA construct of claim 20, wherein said inducible promoter is controlled by a viral protein or an oncoprotein.
22. A pharmaceutical composition comprising at least one of the DNA sequences of any one of claims 17-21 in admixture with a pharmaceutically acceptable carrier.
23. A method of inhibiting papillomavirus associated diseases or cancer comprising administering said pharmaceutical composition of claim 22 to a patient afflicted with a papillomavirus associated disease or cancer.
24. The method of claim 22, wherein said inducible promoter is controlled by the expression of a papillomavirus protein or an oncoprotein in said patient as a result of said disease or cancer.

25. A DNA sequence comprising the full length AAV genome modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 17 to 21.

26. A DNA sequence comprising at least the minimum portion of the AAV genome sufficient to complement a defective rAAV, wherein said portion of the AAV genome is modified to replace the AAV Rep78 DNA sequence with said DNA sequence encoding said AAV Rep78 mutant of any one of claims 17 to 21.

27. A method of producing recombinant AAV (rAAV) comprising transducing a susceptible mammalian cell with a DNA sequence of claims 25 or 26 and a DNA sequence encoding said rAAV to obtain rAAV virions.

28. The method of claim 27, wherein said DNA sequence is sufficient to complement a defective rAAV comprises at least the AAV *lip-cap* gene and said DNA sequence encoding an AAV Rep78 mutant.

29. The method of claim 28, wherein said rAAV comprises a DNA sequence encoding a heterologous gene.

30. A method of treating papillomavirus associated diseases or cancer comprising administering a pharmaceutical composition comprising a AAV Rep78 protein to a patient afflicted with a papillomavirus associated disease or cancer, wherein said AAV Rep78 protein binds to a papillomavirus promoter that controls the expression of a papillomavirus oncoprotein or binds to a promoter that controls the expression of an oncoprotein.

31. The method of claim 30, wherein said AAV Rep78 binds to nucleotides 14-56 of p97 of HPV-16 inhibiting expression of HPV.

32. An AAV Rep78 regulation element comprising the nucleotides shown in the nucleotide sequence of Figure 2, wherein said element provides the binding site for the AAV Rep78 protein.

33. The AAV Rep78 regulation element of claim 32 comprising about nucleotides 14 -56 of the nucleotide sequence of Figure 2.

34. An AAV Rep78 regulatable promoter comprising the regulation element of claims 32 or 33 and the remaining promoter sequences from a promoter other than the HPV-16 p97 promoter.

35. A method of selecting an AAV Rep78 mutant that possesses different biochemical and biological functions as compared to the wild-type AAV Rep78 protein, wherein said method comprises:

contacting an AAV Rep78 mutant of any one of claims 1-14 with at least one DNA sequence obtained from one or more of a papillomavirus, an AAV, a HIV or an oncogene for a period of time to allow binding of said mutant to said DNA; and

determining said binding of said mutant to select said mutant that binds differently to said DNA than said wild-type Rep78 protein.

36. Use of the pharmaceutical composition of claims 15 or 22 for the treatment of a papillomavirus-associated disease, cancer or a HIV-associated diseases.

37. Use of the AAV Rep78 mutant or the AAV Rep78 protein of any one of claims 1-14 or the DNA sequence encoding at least one AAV Rep78 mutant, AAV Rep78 protein or a fusion protein according to any one of claims 1-14 to prepare a pharmaceutical composition for the treatment of a papillomavirus-associated diseases, cancer or HIV-associated diseases